

**IN THE UNITED STATES DISTRICT COURT
FOR THE NORTHERN DISTRICT OF OKLAHOMA**

STATE OF OKLAHOMA, ex rel.)
W.A. DREW EDMONDSON, in his)
capacity as ATTORNEY GENERAL OF)
THE STATE OF OKLAHOMA and)
OKLAHOMA SECRETARY OF THE)
ENVIRONMENT C. MILES TOLBERT,)
in his capacity as the TRUSTEE FOR)
NATURAL RESOURCES FOR THE)
STATE OF OKLAHOMA)

Plaintiffs,)

vs.)

Case No. 4:05-cv-00329-GKF-SAJ

1. TYSON FOODS, INC.,)
2. TYSON POULTRY, INC.,)
3. TYSON CHICKEN, INC.,)
4. COBB-VANTRESS, INC.,)
5. AVIAGEN, INC.,)
6. CAL-MAINE FOODS, INC.,)
7. CAL-MAINE FARMS, INC.,)
8. CARGILL, INC.,)
9. CARGILL TURKEY)
 PRODUCTION, LLC.,)
10. GEORGE'S, INC.,)
11. GEORGE'S FARMS, INC.,)
12. PETERSON FARMS, INC.,)
13. SIMMONS FOODS, INC., and)
14. WILLOW BROOK FOODS, INC.,)

Defendants.)

EXPERT REPORT OF DR. CHRISTOPHER M. TEAF

Qualifications & Experience

1. My name is Dr. Christopher M. Teaf. I am over 18 years of age and am competent to testify. All opinions presented in this statement reflect personal knowledge based on information and data that I have reviewed in this case. All

opinions presented in this affidavit are given to a reasonable degree of scientific certainty.

2. My educational background includes a Bachelor's degree in Biology (with Honors) from Pennsylvania State University (1975), a Master's degree in Biological Science from Florida State University (1980), and a Ph.D. in Toxicology from the University of Arkansas for Medical Sciences (1985), where I conducted my research at the Division of Genetic Toxicology, National Center for Toxicological Research (Jefferson, Arkansas).

3. I presently hold positions as Associate Director at the Center for Biomedical & Toxicological Research and Waste Management at Florida State University (since 1983), as well as Director of Toxicology for Hazardous Substance & Waste Management Research, Inc. since 1985 (President since 1989). I have held adjunct teaching appointments at the Florida State University/State University System Program in Medical Sciences and the Florida A & M University College of Pharmacy and Pharmaceutical Sciences.

4. My ongoing research and scientific advisory interests principally are in the area of health risk assessment for human exposure to occupational and/or environmental chemical, physical and biological hazards. My principal activities for approximately 25 years have included the performance of risk assessments concerning human health and the evaluation of adverse effects of chemical, physical and biological exposures under the requirements of the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA, Superfund), Superfund Amendments and Reauthorization Act (SARA), the Resource Conservation and Recovery Act (RCRA), the Toxic Substances Control Act (TSCA), the Occupational Safety and Health Act (OSHA), and other related federal or state legislation. These professional activities over three decades have involved the evaluation of potential human health impacts of many organic substances (*e.g., petroleum products, pesticides, chlorinated and non-chlorinated chemicals*), as well as inorganic and/or physical agents (*e.g., acids/caustics, metals, particulates, fibers*), and

microbiota (*e.g., molds, fungi, bacteria*) in air, water, soils sediments and other environmental media.

5. For over two decades, I have served as a peer reviewer for publications submitted to numerous scientific journals, including Human & Ecological Risk Assessment, Environmental Forensics, Environmental Toxicology & Chemistry, Chemosphere, Bulletin of Marine Science, Ohio Journal of Science, Journal of Phytoremediation, and Environmental Biology of Fishes, as well as for reviews of research submitted to the Agency for Toxic Substances and Disease Registry (ATSDR) and the World Health Organization (WHO). I presently serve on the Editorial Boards for several of these scientific journals, and am the Senior Human Health Editor for Human & Ecological Risk Assessment, an international journal. I have published many scientific papers, articles and book chapters concerning toxicological effects and risk evaluations related to occupational and environmental exposures and effects. A complete copy of my Curriculum Vitae, including publications, is included as Attachment B to this report.

6. For approximately 25 years, I have directed and conducted research projects and human health education activities for agencies such as the World Health Organization (WHO), the North Atlantic Treaty Organization (NATO), the U.S. Environmental Protection Agency (U.S. EPA), the U.S. Department of Energy (U.S. DOE), the U.S. Department of Agriculture (U.S. DOA), the federal Agency for Toxic Substances and Disease Registry (ATSDR), the Florida Department of Environmental Protection (FDEP), the Florida Department of Health (FDOH), and the Florida Department of Community Affairs (FDCA).

7. I served as Toxicologist representative to the Toxic Substances Advisory Council for the Florida Department of Labor and Employment Security, the agency responsible for implementation of the Florida Right-to-Know Law, from 1990 to 2000, and I was Chair of that group from 1992 to 1996. I serve presently as a member of the Advisory Board of the Center for Terrorism and Public Health within the Florida State University College of Medicine. I served as toxicologist to the Florida Governor's Financial and

Technical Advisory Committee concerning petroleum site remediation issues from 1986-1992, and also as toxicologist representative to the Florida Landfill Technical Advisory Group (1994-1995) for FDEP. I presently serve as Toxicologist representative to the state Petroleum Technical Advisory Committee. In addition, for nearly 20 years I have been toxicologist representative, appointed by the Secretary of the FDCA, and have served for a period of time as Vice-Chairman of the District 2 Local Emergency Planning Committee under the State Hazardous Materials Emergency Response Commission. I was the appointed liaison for the State University System Toxicological Research Coordinating Committee and the (then) Florida Department of Environmental Regulation (FDER) from 1986-1989.

8. Since the late 1970's, I have organized and participated in many full length graduate and undergraduate level academic courses, multi-day technical training courses and seminars for organizations such as the NATO, the WHO, the American Bar Association, U.S. EPA, U.S. DOA, FDOH, Florida Engineering Society, Florida Hospital Association, FDEP, State Fire Marshal's Office, National Hazardous Materials Training Center, Florida State University, University of Florida, Florida A&M University, Georgia Institute of Technology, and the ATSDR. These technical courses and educational activities have included: issues of occupational/environmental chemical or biological exposures and public health (including toxicology and effects of occupational or environmental exposures), risk assessment, risk management, the implementation of health advisories for physicians, and ongoing health concerns regarding environmental contamination and occupational exposure situations in Eastern Europe (*e.g., Bulgaria, Czech Republic, Hungary, Poland*), in Central Asia (*e.g., Kazakhstan*), and in Russia. I have organized, coordinated and taught many 40 hour and 8 hour training courses on health and chemical safety issues related to OSHA requirements for the public and private sectors.

9. I have provided advisory services on toxicology, risk assessment, and health or environmental issues to the U.S. Attorney, the Florida State Attorney's Office and the Attorneys General for Florida, Washington, and Oklahoma. I have been qualified to provide expert testimony to federal and state courts, to administrative proceedings, and

to regulatory or legislative groups concerning toxicology, environmental chemistry, occupational exposures, human health effects of chemical substances, and health risk assessment since 1986.

Retention and Preparation

10. I have been retained in this case by the State of Oklahoma to provide technical analysis and opinions in the areas of toxicology and human health for issues associated with the land application of poultry waste within the Illinois River Watershed (IRW). Specifically, this report addresses health hazards related to bacteria and indicator organisms, to disinfection byproducts, including trihalomethanes (THMs) and haloacetic acids (HAA5s), and to cyanobacteria (also known as blue-green algae or BGA). In preparing my opinions in this case, I have reviewed many documents, reports, datasets, and scientific literature. Materials that I have reviewed are cited herein as appropriate or are included in my formal document production in this matter.

Impairments and Health Hazards in the Illinois River Watershed (IRW)

11. There are demonstrated historical and ongoing hazards and impairments associated with surface water and groundwater in the IRW. These include adverse conditions caused by “pathogens” (e.g., bacteria, viruses, and protozoa as included in the definition and application of the “indicator organisms” element of water quality criteria for Oklahoma and many other states), harmful blue-green algae (cyanobacteria), and chemical disinfection byproducts (i.e., trihalomethanes [THMs] and haloacetic acids [HAA5s]). The Illinois River and most of its tributaries, which include the Baron Fork and Flint Creek among others, historically have been classified by the State of Oklahoma as outstanding water resources for purposes of recreation, wildlife propagation and aesthetic values (OCC, 2004). These hazards and impairments that are present in numerous Illinois River Watershed streams demonstrate conditions that limit the designated uses of the water resources, and they represent an imminent and substantial endangerment to human health.

12. In 2003, the Oklahoma Secretary of the Environment issued a formal report stating that the general health and the water quality of the Illinois River were endangered by pollution from stormwater that runs off from poultry operations or from fields on which waste has been spread (OSE, 2003). Additionally, Engel (2008) developed an approximate poultry waste generation (tonnage) amount for each of the Defendants on a monthly basis, an annual basis, as well as the entire time frame of 1998-2006. His conclusions were that the vast majority of poultry waste generated inside the IRW is disposed inside the IRW, and that a significant amount of poultry waste generated outside of the IRW is disposed inside the IRW as well. Fisher (2008) has reviewed the geology and hydrology of the IRW and has concluded that the IRW is an area of high permeability and that groundwater is particularly susceptible to impact by surface contamination, including land-applied poultry waste. He also has provided estimates of the timing of poultry waste application during the year. Harwood (2008) has addressed a number of critical issues related to bacterial occurrence, origin and significance, concluding that poultry waste poses significant health issues for river users in addition to posing serious environmental problems for the entire IRW. Through various matrix samples, Harwood identified a unique bacterial biomarker which indicates that IRW contamination can be linked specifically to poultry operations and poultry waste land disposal practices.

13. Olsen (2008) has addressed a variety of chemistry and data collection issues to characterize impacts to the IRW, and has demonstrated by evaluating metals, nutrients, other analytes, physical measurements, and indicator organisms, a unique poultry-related chemical "signature". Through his analyses, he concluded that the "signature" has been and is present in environmental samples collected throughout the IRW. Olsen makes the additional distinction that the unique poultry "signature" in surface water is not found in the absence of poultry operations. In addition, statistical measures of common presence for a variety of analytes and microbiota (e.g., Principal Component Analysis or PCA) have been addressed by Olsen (2008). Those measures support a conclusion that the bacterial impacts are related to, and occur coincident with, land application of poultry waste, including manure and litter (CDM, 2008).

14. Cooke and Welch (2008) have reviewed extensive data from the IRW and from Lake Tenkiller, a drinking water reservoir and recreational resource, concluding that the production of algal biomass in Tenkiller Reservoir (also known as Lake Tenkiller and Tenkiller Ferry Lake) is significantly correlated with the changes in total phosphorous and that the mean eutrophic state has ranged from mild to very eutrophic in recent years. The eutrophic conditions have bred and continue to breed blue-green algae which, in addition to causing nuisance tastes, odors, and toxins in potable water, contribute to the production of potentially carcinogenic disinfection by-products during the treatment process of potable water supplies. Caneday (2008) has evaluated recreational usage of the Illinois River and its tributaries, and has provided estimates of users and temporal distribution of recreational activities during the year. Stevenson (2008) has evaluated nutrient concentrations in the IRW and determined the concentrations to be higher than many other regions, as well as being directly related to poultry house densities located in the watershed. He also concluded that nuisance algal biomass as well as modifications in aquatic species composition were related to poultry house densities, coupled with Total Phosphorous (TP) concentrations.

15. The following sections provide information which illustrates why the observed hazards and impairments can be and are harmful to human health, and how those hazards and impairments are linked to the process of land spreading of poultry waste in the Illinois River Watershed.

Bacteria

16. There are currently nine water bodies within the Oklahoma portion of the IRW that are identified by the Oklahoma Department of Environmental Quality (ODEQ) and listed on the 2006 U.S. EPA 303(d) list as “impaired” for the beneficial use defined as Primary Body Contact Recreation (PBCR; Table B1, Figure B1) as a result of bacteria and indicator organisms, including *E. coli*, enterococci, and/or the fecal coliforms (ODEQ, 2006a). It should be noted that one segment (OK 121700030080) of the Illinois River inadvertently was omitted from the 2006 Oklahoma U.S. EPA 303(d) list comprised of 8 segments, an oversight which subsequently has been corrected (ODEQ, 2008). That omission represented a 31.5-mile stretch on the main stem of the Illinois River where a

large amount of intensive aquatic recreational activity on the river takes place (Caneday, 2008). With the inclusion of that 31.5-mile segment, over 75% of the main stem of the Illinois River and its major tributaries is considered to be “impaired” for the category of “pathogens” (Figure B1), due to the presence of fecal indicator organisms, including bacteria.

17. The U.S. EPA 303(d) List is prepared on a biennial basis as an ongoing obligation under the 1972 amendments to the Clean Water Act and requires states to compile a list of water bodies that are “impaired” for various parameters, and to submit updated lists of the impaired water bodies to the U.S. EPA every two years (ODEQ, 2006a). The U.S. EPA 303(d) list defines an “impaired” water body as one which does not meet the state water quality standards, in this case the applicable standards are those related to Primary Body Contact Recreation with surface water. The presence and magnitude of microbial indicator organisms, as used in the context of impairment determinations, are commonly used and widely accepted measures of the potential for presence and health significance of pathogens, including bacteria, viruses and protozoa (Toranzos et al., 2002; WHO, 2003; NRC, 2004; EPA, 2005; Wade et al., 2006).

18. For the Oklahoma portion of the Illinois River’s main stem and its major tributaries, approximately 77% is listed on the EPA’s 2006 303(d) list. With respect to individual segments, the entire lengths (100%) of Sager, Tyner, Peacheater, and Flint Creek are designated as impaired. The Baron Fork is 66% impaired and Tahlequah Creek is 77% impaired (Figure B1). Caney Creek was taken off the 2006 303(d) list; however, water samples from the IRW segment, still show regular exceedances of indicator organisms, for which that water body failed several pathogen criteria in 2004 and 2002. It is important to note, however, that segments which are listed on the 303(d) impaired list, are not the only surface water sites within the IRW with exceedances of health-based water criteria. In addition, multiple detections of fecal indicator organisms have been found in Edge-of-Field samples, soils and sediments samples and other creeks/streams located in the IRW.

19. Throughout the Oklahoma portion of the IRW, land use designation primarily falls into one of two categories: forest or pasture (EPA, 2001a). Urban impacts are minimal in the Oklahoma portion of the watershed, with only three permitted National Pollutant Discharge Elimination System (NPDES) locations present, coupled with low human population densities in the Oklahoma portion of the IRW (EPA, 2007b). All three of these locations are present either far down on the Illinois River (i.e., Tahlequah) or they are located on downstream tributaries (e.g., Baron Fork and Caney Creek). Their impacts are further limited due to the relatively low permitted discharge rates, ranging from 0.28 to 5.27 million gallons per day (MGD; Table B3). Sewage bypasses and overflows are not significant sources of contamination under normal conditions. The Waste Water Treatment Plant's (WWTPs) in the IRW are all separate sewage systems (SSS's) which do not combine untreated waste with stormwater runoff, thus should not be as heavily impacted by large rain events as a combined sewage system (CSS) would be. Finally, the permitted bacterial limits are restrictive, thus controlling the total amount of bacteria that can be released from the systems.

20. Contamination of surface water and groundwater supplies by bacteria has long been recognized as a human health concern in the United States and around the world. The 1986 U.S. EPA *Ambient Water Quality Criteria for Bacteria* provided historical context and recommendations concerning appropriate guidelines for microorganisms (U.S. EPA, 1986). Subsequent refinements and updates to that guidance are represented by the *Implementation Guidance for Ambient Water Quality Criteria for Bacteria* (U.S. EPA, 2003). This health-based guidance fits into the operable "fishable/swimmable" goals of the Clean Water Act, which specifically requires that water quality standards must "protect the public health and welfare, enhance the quality of water, and serve the purposes of this Act."

21. Microbiological contamination of water can be caused by bacteria, viruses, protozoa and other related organisms. The number and diversity of these potential contaminants has resulted in the development of practical assessment and protection strategies which employ "indicator organisms" as a surrogate for the quantification of specific species in water bodies (Barrell et al., 2000; U.S. EPA, 2003; National Research

Council, 2004). These indicator organisms, such as *Escherichia coli* (*E. coli*), enterococci, and fecal coliform bacteria, may not cause illness directly, but they have demonstrated characteristics which make them reliable indicators of other harmful pathogens in water (Wade, 2006). Although the most commonly reported illnesses associated with bathing in contaminated water typically are gastrointestinal in nature, other illnesses and conditions affecting the eyes, ears, skin and upper respiratory tract can occur as well. Essentially all local, state, and national health agencies employ one or more of the indicator organisms in their water quality management programs, and this is true internationally as well (WHO, 2000; EPA, 2003). Thus, there is consensus that the presence of these indicator organisms at levels greater than the health-based criteria or standards represents a human health threat.

22. The 2003 U.S. EPA *Implementation Guidance* cited previously, provided detailed information regarding the basis for the environment and health agency recommendations, including discussions on the epidemiology of microbiological disease related to water uses such as swimming, kayaking, water skiing, and other activities where direct contact and immersion in the water are likely. For *E. coli*, a geometric mean density of 126 organisms per 100 milliliters (ml) of water over a 30-day period was associated with an illness rate of 0.8%, or 8 illnesses per 1,000 recreational users. As a short-term measure, this 0.8% illness rate was associated with bacterial counts of 236 per 100 ml as an upper limit. For the enterococci, a geometric mean of 33 organisms per 100 ml and an upper limit of 62 organisms per 100 ml were associated with the 0.8% illness rate (OAC, 2007). Above these threshold levels, the agency noted that illness rates rise sharply, and the health-based recommendation seeks to remain below that part of the statistical curve (Figure B2). The State of Oklahoma, along with many other states, has adopted these indicator organism criteria as a fundamental element of their water quality criteria for protection of human health.

23. The exposures represented by recreational uses of the Illinois River and its tributaries, as described by Caneday (2008), both floating and non-floating, are consistent with the types of activities considered by the EPA (2003) in its estimates of illness rates of 8 per 1000 people (0.8%) for indicator organisms in fresh water. Using

the figures for intensive uses of the Illinois River Watershed provided by Caneday (2008), it is possible to estimate that over 1,200 illnesses are occurring on an annual basis even if the bacterial water quality is just being met. While states can determine their own disease incidence targets, the EPA (2003) recommends less than 1.0% illness rate (e.g., 0.8% is used by many states). This is in part because the epidemiological data illustrate an exponential increase in rates beyond the 1.0% illness incidence threshold, instead of a linear increase. Stated differently, at low indicator organism densities (i.e. below the health-based recommendation) the pattern of increase in illness is fairly small and the line is flat, while at higher indicator organisms densities, the illness rate curve rises much more steeply (Figure B2).

24. Primary Body Contact Recreation (PBCR) is an exposure category defined by the State of Oklahoma Administrative Code in Title 785, Chapter 45 (OAC, 2007). This beneficial use category is specifically intended to protect recreational users of water bodies from contamination during the recreational season of May 1 through September 30. PBCR is defined as involving direct body contact (i.e., the dermal exposure route) with surface water through activities such as swimming, wading, canoeing, and fishing. It should be noted that PBCR also explicitly includes the additional likelihood of incidental water ingestion (i.e., oral exposure route) while recreation is occurring in the watershed through multiple activities (OAC, 2007). In water bodies that are governed by the PBCR requirements, the State of Oklahoma mandates that such water “shall not contain chemical, physical or biological substances in concentrations that are irritating to skin or sense organs or are toxic or cause illness upon ingestion by human beings” (OAC, 2007). Clearly, levels of bacteria and indicator organisms that exceed health-based criteria and other standards will and do pose such an unacceptable health risk to users of the Illinois River and its tributaries.

25. Information that is available regarding land disposal of poultry waste shows that nearly two-thirds (over 63%) of waste spreading occurs in the months of February through June (Figure B3), based upon data for the period 1999-2004 (Fisher, 2008; Engel, 2008). Information available regarding the pattern of a major PBCR use (“floating”) within the Illinois River and its tributaries in the 2004-2007 period by month shows that

99% of that usage type occurs in the months of May through September (Caneday, 2008; Figure B3). Floating includes ancillary direct contact activities in the water such as swimming and wading as well, and reasonably can be considered to include an ingestion component of exposure, as well as dermal. Thus, a very large proportion of total annual land spreading of poultry waste is conducted during the months which just precede or overlap with the times of maximum direct contact recreational use of the Illinois River and its tributaries. Coincidentally, the rainy season in Oklahoma tends to be between February and June (NOAA, 2000), which coincides with and immediately follows the highest rates of waste application (Adamski, 1987; Adamski & Steele, 1988; Fisher, 2008). The rainy season also immediately precedes the highest rates of floating activity (Figure B4). On average, approximately 45% of annual rainfall in Northeast Oklahoma occurs between the months of February and June (NOAA, 2000). Rainfall, specifically when it occurs shortly after land spreading, results in microbial pathogen distribution as a result of runoff from spread poultry waste or by leaching through the soil profile (Giddens and Barnett, 1980; Gagliardi and Karns, 2000; Fisher, 2008; Olsen, 2008), even if buffer zones are used correctly, which they frequently are not.

26. Recently, over six consecutive years (2000-2005), there has been widespread measurement of the presence of indicator organisms in surface waters of the IRW, which indicate that those waters broadly and regularly exceed the Oklahoma Water Quality Standards and/or health-based screening levels (OWRB, 2007; CDM, 2008). Single point criteria for surface water along the impaired water body segments from recreational seasons 2005-2007 were exceeded in a large number of samples. Most of these samples are at least twice the standard and the values may exhibit exceedances up to a maximum of 125 times the standard (Table B2). Another example of regular and widespread exceedances of particular interest exist in specifically identified public access (PA) locations (Figure B5), most of which are within impaired water body segments, where a remarkable 13 of 15 locations (nearly 90%) had results where one or more indicators were in excess of the Oklahoma standards for fecal bacteria as shown by indicator organisms (CDM, 2008). In addition to single point criteria, a number of multi-sample geometric means within impaired water body segments, according to the

procedures of the Oklahoma Water Resources Board, also showed exceedances. The State of Oklahoma mandates that a geometric mean calculation for Primary Body Contact Recreation have no less than five samples for any 30-day time frame of interest (OAC, 2007). Results for the recreational seasons of 2005-2007 show exceedances in all water bodies where sufficient samples were taken to perform the calculation, with those exceedances being in the range of 2 to 30 times the standard (Table B2). While Peacheater Creek, Tyner Creek, Tahlequah Creek, and a portion of the Illinois River did not have adequate sampling frequency within the specified time frame to fairly calculate the geometric mean, the percentage of single point criterion exceeding the standard was in most cases nearly 20% of the samples, at levels that were 2 to 65 times greater than the health-based screening level (CDM, 2008; Table B2). These results, in my opinion, demonstrate a chronic and persistent problem of bacterial contamination which represents an imminent and substantial endangerment.

27. In addition to Primary Body Contact Recreation uses associated with surface water exposures, significant exposures to contaminated groundwater can occur through drinking water supplies as well. In Oklahoma, groundwater is protected such that bacterial levels in groundwater must be “nondetect”, which is reasonable based on the absence of normal bacterial fauna in groundwater at deeper depths (NRC, 2004; EPA, 2006A; USGS, 2007b). Once groundwater pollution has been determined to have occurred via human activities, the water supply is to be restored to a quality sufficient to support its designated beneficial use, in this case potable water supplies (OAC, 2007). Over 1,700 groundwater wells have been identified in the Oklahoma portion of the IRW, of which 98% are used for domestic purposes (OWRB, 2008; Fisher, 2008; Figure B6). Figure B7 illustrates the CDM groundwater well sampling sites within the IRW while Figure B8 shows the locations of Geoprobe samples taken within the IRW. Bacterial contamination of groundwater, including impacts by *E. coli*, fecal coliforms and enterococci, has been demonstrated in many shallow wells and in other wells at depths to approximately 150 feet, as well as in a majority of shallow groundwater Geoprobe samples (CDM, 2008). Olsen (2008) identified the poultry “signature” in the majority of these samples. While these Geoprobe samples are not necessarily

completed to depths from which groundwater is taken for domestic purposes, they clearly demonstrate the potential pathway between microorganisms at the surface and those which can contaminate groundwater. These elements further demonstrate the vulnerability of shallow groundwater (NRC, 2004; Fisher, 2008), and illustrate the health concerns that are associated with land spreading of poultry waste in areas where groundwater is used for potable and other domestic purposes.

28. Bacterial levels of human health significance also have been found in a number of springs within the IRW. At their point of release from the ground, where they “daylight”, springs represent a transition from groundwater to surface water. Thus, they can be indicators of impacts to groundwater and/or surface water (Fisher, 2008). Springs sampled in the Oklahoma portion of the IRW are shown in Figure B9. Approximately 21% of the spring samples exceeded the surface water bacterial standards. Most of those samples were located in close proximity to the Illinois River and its tributaries. Olsen (2008) concluded that nearly 40% of spring samples were impacted by poultry waste.

29. Bacteria have been detected in surface water, groundwater and springs in the IRW at levels that are indicative of significant concern from a human health perspective (CDM, 2008). Edge-of-Field samples for fields that have received recent spreading of poultry waste have shown bacterial colony counts in water which are similar to those reported for water samples into which, raw, untreated sewage (i.e., 10^5 or 100,000 MPN or greater) has been spilled from U.S. sewage treatment plants (Metcalf & Eddy, 1991; Brosnan et al., 1996; CDM, 2008; Harwood et al., 2005). Of the 22 Edge-of-Field sample locations which exhibited bacteria and indicator organism results in excess of 100,000 MPN per 100 milliliters, seven of the locations exhibited results for enterococci, fecal coliforms and/or total coliforms greater than or equal to 1,200,000 MPN per 100 milliliters, with three sites having a total of 5 reports of greater than or equal to 1,600,000 MPN per 100 milliliters (Table B6). Through Principal Component Analysis (PCA), it has been determined that 100% of the Edge-of-Field samples showed poultry related impacts. Furthermore, of the 50 highest “PC1” (poultry) scores of all sampled media, 44 were Edge-of-Field samples, with the highest score resulting in a sample

taken from field runoff immediately after poultry waste application (Olsen, 2008). In addition, the maximum levels for *E. coli*, enterococci and fecal coliform found in poultry litter/waste samples collected by CDM in 2006 were 120,000 MPN per gram of litter (CDM, 2008). Other impacted media include sediments and soils as demonstrated by Fisher (2008) and Olsen (2008).

30. In addition to the numerical comparisons between health-based criteria and detected levels of bacteria and indicator organisms as a measure of potential health hazard, it is useful to consider the relative importance of microbial sources in the IRW as well. Processes to accomplish this have been developed by USEPA and a number of individual states, including Oklahoma, under the Total Maximum Daily Load (TMDL) program (USEPA, 1997; USEPA, 2001b; ODEQ, undated).

31. An analysis of potential sources for fecal coliforms was conducted in a fashion consistent with that employed by USEPA and ODEQ for the six counties which share some portion of the Illinois River Watershed (Adair, Cherokee, Delaware and Sequoyah in OK; Benton, Washington in AR). That analysis considered fecal coliform contributions by a variety of categories for which data were available, including: domestic pets, deer/wildlife, failing septic systems, permitted point sources (i.e., NPDES outfalls), and livestock. The livestock category was further subdivided into groups by poultry, cattle/calves, horses/ponies, sheep/lambs, and swine. Table B4 summarizes the contributions for each source category, and also provides a summary of the relative contribution from the five livestock categories. The numerical values for each category are expressed in units of Colony Forming Units per day (CFU/day). For example, the total fecal coliform load from poultry and from cattle/calves is approximately 5×10^{15} CFU/day, or 5,000 trillion CFU/day each. Table B5 provides the underlying summary calculations and input parameters for the values presented on Table B4.

32. Several important conclusions can be drawn from this source contribution analysis, including the following:

- The categories of domestic pets, deer/wildlife, failing septic systems and point sources each contribute from 0.01% to 0.9% of total fecal coliform loading. Those contributions are not significant in comparison to the contribution from livestock;
- The livestock category alone contributes nearly 99% of total fecal coliform loading;
- Within the livestock category, poultry and cattle/calves each contribute just over 40% each of the total, swine contribute about 14% of the total, sheep/lambs contribute about 0.1% of the total, and horses/ponies contribute about 0.03% of total fecal coliform loading;

In addition, leachability of poultry waste was on the order of 1 to 5 times higher than fresh cattle manure, and is likely to be even greater for dry manure based on the smaller particle sizes present in poultry waste (Olsen, 2008). Therefore, poultry waste is much more likely to leach components with the potential for adverse impacts from the site of application to nearby water sources, than is cattle manure.

33. For over a decade, the practice of adding antibiotics (e.g., fluoroquinolones, tetracyclines, aminoglycosides, macrolides) to farm feed has been widely used to fight microbial infection in poultry and other livestock, as well as sub-therapeutically to increase feed conversion efficiency and weight gain (Nandi et al., 2004; PCIFAP, 2008). Concern has been raised about the ability of bacteria such as *E. coli*, *Salmonella*, *Campylobacter*, and *Staphylococcus*, which are common in poultry, to become antibiotic-resistant, thereby representing a source of infection in humans as a result of this widespread antibiotic use in farming (White et al., 2003; USFDA, 2002; Nandi et al., 2004; Hurd et al., 2004; Diarra et al., 2007; Dupont, 2007; Smith et al., 2007; PCIFAP, 2008). Price et al. (2007) reported that individuals who work in the poultry industry were much more likely to carry antibiotic resistant strains of *E. coli* than other members of the community. According to the World Health Organization, the “use of antimicrobials outside of human use is of serious concern given the alarming emergence of bacteria, which have acquired, through this use, resistance to antimicrobials” (WHO, 2008a). These concerns ultimately led the European Community to prohibit the use of antibiotics as growth promoters. Antibiotics used by the poultry industry, including defendants in this case, to treat or control bacterial diseases include: Baytril

(enrofloxacin), Sarafloxacin, Bacitracin (BMD), Penicillin (e.g., Ampicillin, Amoxicillin, Methicillin), Gentamycin sulfate, and tetracyclines (e.g., chlortetracycline (CTC), neoterramycin, oxytetracycline), among others (see e.g., Bates #: TSN088218SOK, TSN088077SOK, TSN088197SOK, CM003570, SIM AG09496, CARTP109186). In addition to antibiotics, pesticides (e.g., Larvadex, Fenbendazole, Piperazine, Levamasol) are also added to feed to control worms and fly larvae (see e.g., Bates #: CM003473, CARTP169958, CARTP143292, CARTP158787), and thus would be present in excreted poultry waste. This information clearly illustrates that a number of significant pathogens are a recognized and inherent health concern associated with poultry raising operations, and that the industry historically has actively treated the flocks to control these very bacteria in poultry waste which are potentially dangerous to the flocks.

34. It is evident that the poultry industry is and has been aware for some time of the potential health problems that these pathogens represent due to the widespread use of the veterinary antibiotics (see previous paragraph). Before the use of fluoroquinolones became common in animal husbandry (including poultry), antibiotic-resistant bacterial strains were virtually nonexistent (Bren, 2001; WHO, 2008b). In reference to the benefit of fluoroquinolone use being compromised, Hofacre et al. (1999) noted that the "Use of antibiotics in both humans and animals contributes to the selection pressure resulting in this resistance". *Campylobacter* from 28% of human patients in one study showed resistance to fluoroquinolones (Ellis-Pegler et al., 1995). In one particular case, *Salmonella minnesota* and *E. coli*, isolated from a number of turkey poult, were subjected to an antibiotic susceptibility pattern test. In that analysis, *E. coli* showed resistance to 13 of the 16 antibiotics tested and *S. minnesota* showed resistance to 10 of the 16 antibiotics tested (University of Missouri Veterinary Lab, 2007). It appears that the turkeys in question must have been subjected to numerous antibiotics in feeding operations, judging by their resistance to such a broad list of antibiotics. However, this particular case simply further illustrates what White et al. (2000) previously had demonstrated: that antibiotic-resistance is and has been on the rise. *Campylobacter* have been shown elsewhere to be broadly resistant to macrolide antibiotics (Bolenger and Shryock, 2007). In their study, White et al. (2000) showed avian (poultry) pathogenic *E.*

coli (APEC) resistance to Sarafloxacin increased from 15% in 1996 to 40% in 1999 and dual-resistance to Sarafloxacin and to Enrofloxacin increased from 9% in 1997 to 30% in 1999. To further illustrate the growth of multi-antibiotic resistance over time, three separate studies performed by Bass et al. (1999), White et al. (2000), and Zhao et al. (2005) published APEC results of 64%, 66%, and 71% of samples, respectively, having resistance to five or more antibiotics. With respect to individual antibiotics, the same three studies demonstrated that the use of tetracyclines has become so extensive that APEC-resistant strains ranged from 85 to 89% of bacteria, while Gentamycin resistance is in the range of 62 to 69% of APEC strains. Therefore, the spreading of poultry waste material clearly aids in the dispersion of antibiotic resistance in the environment, creating imminent and substantial endangerment to those using the river, as well as contributing to this recognized worldwide problem of antibiotic resistance.

35. Land spreading of poultry waste has long been recognized as a major bacterial contamination source (Crane et al., 1980; Adamski, 1987; Adamski and Steele, 1988; PCIFAP, 2008). Spreading of waste material, a traditional agricultural waste disposal practice, becomes a major source of contamination because frequently it exceeds the rate at which wastes can be accommodated by or processed in agricultural ecosystems (Coyne and Blevins, 1995). Rainfall, specifically when it occurs shortly after land spreading, may then result in pathogen distribution by runoff from spread poultry waste or by leaching through the soil profile (Giddens and Barnett, 1980; Gagliardi and Karns, 2000; Fisher, 2008; Olsen, 2008), even if buffer zones are used correctly, which they frequently are not. This is rendered even more important by the fact that the recreational season for the IRW overlaps with and immediately follows the rainy season, a period which is well within the survivability duration of the bacteria in question (Figure B4). Furthermore, the leachability of poultry waste components into groundwater significantly outweighs that of cow manure (Olsen, 2008). The environmental survivability of bacteria can be on the order of several days to many months (Jamieson et. al., 2002; Tetra Tech, 2004; Davis et al. 2005). Runoff from waste-spread fields carries excess nutrients, pollutants, and pathogens to nearby waterways, which negatively affects surface water, groundwater, aquatic life, and human health;

even months after land application of waste, fecal coliforms and *E. coli* can be resuspended from sediments and transported downstream (Coyne and Blevins, 1995; Hartel et al., 2000; Davis et al., 2005; Ringbauer et al., 2006).

36. Bacteria of human health significance, including *Campylobacter*, *Salmonella*, *Staphylococcus*, *Escherichia coli* and other important species, as well as bacterial indicator organisms such as fecal coliforms and enterococci, are present in poultry waste (Kelley et al., 1995; Jenkins et al., 2006; CDM, 2008; PCIFAP, 2008). The presence of microbial indicator organisms in surface and groundwater bodies suggests that other dangerous bacteria such as *Campylobacter*, *Salmonella* and/or *Staphylococcus* also may be present, in addition to ancillary viruses and protozoa that are more difficult to monitor (e.g., *Cryptosporidium*).

37. *Campylobacter* is a common intestinal bacterium found in a wide range of poultry, domestic livestock, and wildlife (Neill et al., 1984; Lindblom et al., 1986; Kazwala et al., 1990; Waage et al., 1999; Chen et al., 2006; Doyle and Erickson, 2006), though it is quite commonly associated with poultry operations and products globally (Hald et al., 2007), and it is more prevalent in poultry than in swine or cattle (Belanger and Shryock, 2007). While often referred to primarily as a foodborne illness, *Campylobacter* may be waterborne as well (Allos, 2001; O'Reilly et al., 2007). The major effects and complications of *Campylobacter* infection can occur in or near the gastrointestinal tract (USDA, 1991; Allos, 2001; Murray et al., 2003); however enteric (intestinal) diseases are not the only cause for concern. Human campylobacteriosis frequently presents as a sporadic infection, such that it is common to find individual cases in contrast to related outbreaks affecting a large group (Friedman et al., 2000). This sporadic occurrence pattern and the frequently self-limiting nature of the infection causes campylobacteriosis, as well as other enteric diseases that are transmitted by the waterborne route, often to be underreported to public health agencies (Belanger and Shryock, 2007; Leclerc, 2002). Even when diagnosed, campylobacteriosis is reported infrequently (Allos, 2001), and Mead et al. (1999) concluded that less than 3% of diagnosed *Campylobacter* gastroenteritis cases are reported to health authorities.

38. *Salmonella* sp. are bacteria also frequently associated with poultry (Barbour and Nabbut, 1981; Jacobs-Reitsma et al., 1994; Himathongkham et al., 1999; Doyle and Erickson, 2006; Li et al., 2007), and numerous *Staphylococcus* species have been isolated from chickens, including *S. aureus*, *S. sciuri*, *S. gallinarum*, *S. lentus*, *S. chnii*, *S. xylosus*, and *S. warneri* (Adegoke, 1986; Terzich et al., 2000). Infectious diseases related to fecal bacteria from poultry waste include campylobacteriosis, giardiasis, cryptosporidiosis, salmonellosis, *E. coli* 0157:H7, and others (Doyle and Schoeni, 1987; Leclerc et al., 2002; Dipineto et al., 2006; Doane et al., 2007; Dupont, 2007; OSDOH, 2007; Trampel et al., 2007). The State has recognized the potential impacts of chicken waste under such acts as the Oklahoma Confined Animal Feeding Operation and Poultry Registration Acts (ODAFF, 2008). Recognition of human disease acquired by animal sources by the Experts Scientific Workshop has resulted in placing the highest priority for further research on contamination from poultry and other agricultural animals (U.S. EPA, 2007c). Gastroenteritis, caused by fecal pathogens and characterized by nausea, cramps, diarrhea, and vomiting, is a common and serious affliction associated with waterborne bacteria (OSDOH, 2007; VGHI, 2007). Following the review of 22 articles, Pruss (1998) concluded there to be a causal relationship between gastroenteritis and recreational water quality, as measured by bacterial indicators. Important infections caused by *Staphylococcus*, including recent antibiotic resistant strains (e.g., Methicillin-resistant *Staphylococcus aureus*, MRSA), involve skin, and may result in boils, cellulitis, impetigo and serious wound infections. Many other organ systems in the human body can be affected as well (Kloos and Bannerman, 1999; Murray et al., 2003). Strains of *Salmonella* and other bacteria have undergone adaptations to become antibiotic resistant, as described previously in the report.

39. The Oklahoma State Department of Health (ODOH) maintains statistics of specific reportable diseases including diseases caused by bacteria such as *Campylobacter*, *Salmonella*, and *E. coli* 0157:H7, and microscopic parasites such as *Giardia* and *Cryptosporidium*. As described elsewhere in this report, these organisms have been associated with poultry waste and often are also associated with contaminated drinking water, fecal material, and contact with birds. An evaluation of OSDOH records for

Oklahoma counties in the IRW shows that Adair County reported rates of campylobacteriosis considerably in excess of the state average for the period 1997 to 2005 (ODOH, 2006; see Figure B10). Adair County makes up the largest portion of land area within the IRW. In addition, rates of salmonellosis reported between 1990-2005 also have periodically exceeded the average statewide incidence rate. The rate of salmonellosis in Sequoyah County was reported to exceed the state rate for all except three years during the period 1990 to 2001 (ODOH, 2006; see Figure B11). Furthermore, the data from the ODOH shows no associations between serotypes of the *Salmonella* bacterium. In addition to the lack of commonality between serotypes, no common relationships between individuals, demographic characteristics, or locations were identified, as would be expected from a single, large food-borne outbreak (ODOH, 2006).

40. While the ODOH has not investigated any “outbreaks” with regard to the diseases discussed above, it cannot be presumed that incidents of infection are not occurring. Without question, tourism within the watershed is extensive, considering for example, that the general population of Adair County during the period 1990 to 2000 ranged from about 19,000 to 21,000 people (U.S. Census, 1990; U.S. Census, 2000) and at least an estimated 155,000 people use the IRW annually (Caneday, 2008). When using the CDC’s guidelines for investigating an outbreak, a clustering of sickness must take place to warrant an investigation. This, therefore, would be very difficult to achieve under recreational use circumstances, recognizing how many tourists visit the Illinois River watershed from Arkansas, Kansas, Missouri, as well as other counties in Oklahoma. Lee et al. (2002) noted that outbreak investigations were increasingly difficult to document when users convene onto one venue and then geographically disperse. This clearly identifies one possible reason why no focused investigations have been initiated for the Illinois River Watershed. Latency periods in the order of a day to a week (Mayo Clinic, 2008; CDC, 2008), depending on the bacterium, would surely affect reporting statistics if recreational users and tourists to the region are taken into account, and consideration is given to the likelihood of returning to their homes after visiting the IRW. Additionally, outbreaks associated with some infective organisms are

less likely to be investigated than acute diseases characterized by short incubation periods, serious illness requiring medical treatment, and those having recognized etiologies (Lee et al., 2002; Blackburn et al., 2004). Individual sensitivity and enhanced susceptibility among groups such as children, the elderly, and the immunocompromised, further complicates the effectiveness and applicability of disease surveillance (WHO, 2003; NRC, 2004).

41. It is clear that many diseases are commonly under-reported, given the limitations of the passive disease surveillance systems presently in place. Multiple factors play a role in whether disease outbreaks are recognized, investigated, and/or reported, which typically will result in under-reporting of the true illness rate (Lee et al., 2002; Blackburn et al., 2004; Liang et al., 2006; Craun & Calderon, 2006;). Multiple studies (Lee et al., 2002; Yoder et al., 2004; Blackburn et al., 2004; Liang et al., 2006) have concluded that the data which are collected most commonly pertain to “outbreaks,” with no mechanism to include seemingly sporadic cases, and therefore the data do not necessarily represent actual endemic trends with waterborne illnesses. The observations already available concerning disease occurrence in northeastern Oklahoma underscore the potential for increases in infectious diseases related to land disposal of poultry waste in large quantities.

42. The bacterial impacts associated with the application of poultry waste to lands in the Illinois River Watershed represent an imminent and substantial endangerment to human health.

Trihalomethanes (THMs) and Haloacetic Acids (HAA5s)

43. One human health risk associated with the spreading of poultry waste on agricultural fields in large quantities, with associated runoff, is related to the formation of potentially carcinogenic substances that may occur in treated drinking water supplies (Cooke and Welch, 2008). Formation of these products can occur during the normal municipal treatment of water prior to distribution, and is a result of increased organic matter in the raw intake water which combines with chlorine to form “disinfection byproducts” (DBP’s). Several examples of which are trihalomethanes (THMs) and

haloacetic acids (HAA5s). One direct and/or indirect source of this increased organic matter is contaminated runoff from fields on which poultry waste have been spread. Studies have shown there to be an increase in the poultry population and thus an increase in the amount of waste generated by poultry dating back as far as 1950 (Engel, 2008; Fisher, 2008). The total amount of waste generated annually within the IRW was estimated to be over 300,000 tons (Fisher, 2008). Increased organic matter, in conjunction with excessive nutrients, primarily nitrogen and phosphorus, from poultry waste are principal recognized causes of eutrophication, algal growth, and water quality degradation (Lee and Jones, 1991; Cooke and Welch, 2008; PEW, 2008). This scenario is illustrated in Figure T1. The principal source of phosphorus to Lake Tenkiller has been attributed to poultry waste applied to pastures that eventually reaches the Illinois River and then Lake Tenkiller (Engel, 2008; Cooke and Welch, 2008). One of the most effective methods to control eutrophication and restore water quality is by reducing the excessive phosphorus sources (Sas et al., 1989; Lee and Jones, 1991; Slaton et al., 2004; Cooke and Welch, 2008), which in this case could be accomplished by decreasing or eliminating the contribution made by poultry waste.

44. Nutrient pollution to surface waters from “insufficiently treated sewage, runoff from fertilized agricultural areas and lawns, manure and more complex effluent from livestock industries” (Briand et al., 2003), can and does result in eutrophication and algal proliferation. The increase in these algal communities, including those of cyanobacteria, disrupt the complex aquatic environment within the system, which can lead to oxygen depletion, taste and odor issues, and the formation of organics in the water (Fruh, 1967; Hoehn et al., 1980; van Steenderen et al., 1988; Cooke and Welch, 2008; Stevenson, 2008). Cyanobacteria, such as *Microcystis aeruginosa*, have been shown to be productive precursors of trihalomethane disinfection byproducts in treated water supplies (van Steenderen et al., 1988).

45. Chlorine treatment, which has been the most widely used and cost effective disinfectant practice for U.S. public or private water supplies since the early-1900’s and which is commonly used in wastewater treatment facilities (NAS, 1987; Bull et al., 1995), may result in the formation of chemicals known as DBPs, which are of ongoing health

concern (Wistrom et al., 1996; USEPA, 2006c). These DBPs include the predominant class of trihalomethanes, or THMs (consisting of chloroform, bromodichloromethane, dibromochloromethane, bromoform), as well as the haloacetic acids, or HAA5s, (consisting of monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, monobromoacetic acid, and dibromoacetic acid). In addition to these groups, other disinfection byproducts are continually being evaluated for occurrence, toxicity and potential human health risk (Krasner et al., 2006). The formation of DBPs in treated water is a well-recognized phenomenon, and depends on factors such as pH, nature/concentration of organic matter, chlorination contact time, water temperature, chlorine dose, chlorine residual, and bromide concentration (Stevens et al., 1976; Reckhow and Singer, 1990; OWRB, 2005; USEPA, 2006b; Gopal, 2007; Cooke and Welch, 2008).

46. The formation of DBPs is correlated significantly with the content of dissolved organics in raw water (Martin et al., 1993; Stepczuk et al., 1998; Stuart et al., 1999; Gallard and Gunten, 2002; Brown, 2003). Under normal circumstances, naturally occurring organic material (NOM) is a principal precursor with which halogens (e.g., chlorine and bromine) can react to form DBPs (Rook, 1976; Oliver and Lawrence, 1979; Marhaba and Kochar, 2000; Gopal et al., 2007; Richardson et al., 2007; Wang et al., 2007). However, nutrient enrichment (e.g., loading of phosphorus, Total Organic Carbon (TOC), and nitrogen) in water bodies from activities such as spreading of poultry waste to pasture lands, can increase organic matter concentrations and dramatically increase or speed up eutrophication processes in lakes, often with accompanying increases in algal growth, which also can result in increased DBPs (Peterson et al., 1993; Preusch et al., 2002; Lee et al., 2007; Cooke and Welch, 2008). Water samples taken directly following waste application and right after rainfall (Edge-of-Field) from the IRW show high levels of TOC.

47. Excess nutrient loading and nutrient pollution in some circumstances can originate from urban runoff, industry, automobiles, and human sewage. However, in many areas the largest source comes from agriculture and from uses of manure produced by livestock, including poultry, on fields (Anderson et al., 2002; Tang et al.,

2005). Engel (2008) concluded that excess amounts of these fertilizer constituents, in the form of manure and litter spread in quantities beyond that which can be sequestered by plants and soil, lead to increased amounts of phosphorus in runoff. In addition, Engel (2008) concluded that close to 80% of the net annual phosphorus contribution to the IRW comes from poultry. High levels of phosphorus in runoff act to fertilize the aquatic life in water, leading to excessive growth of algae and underwater plants that can lead to production of DBP's (Hoehn et al., 1980; Dore et al., 1982; Lee and Jones, 1991). Enhanced nutrient loading can also increase the formation of *Cladophora*, a branching filamentous green alga, often times found on rocky substances. Stevenson (2008) showed that the algal biomass density was related to total phosphorus concentrations and poultry house density along the streams and rivers in the Illinois River Watershed. Other technical literature has shown that *Cladophora* can be a secondary habitat for pathogenic bacteria (Byappanahalli et al., 2003; Whitman et al., 2003; Ishii et al., 2006), which greatly extends the survivability of the bacteria in the environment. When levels of TOC and DOC decrease, THM precursors and THMs levels decrease as well.

48. The U.S. EPA, the International Agency for Research on Cancer (IARC), the National Toxicology Program (NTP) and other researchers have concluded that the weight of evidence from animal and human studies warrants classification of the THMs (chloroform, bromodichloromethane, dibromochloromethane, bromoform), as well as at least two of the haloacetic acids (dichloroacetic acid, trichloroacetic acid) as Probable or Possible human carcinogens (Cantor et al., 1978; Morris et al., 1992; Nokes et al., 1999; USEPA, 2007a; Villanueva et al., 2007). One case-control study from Canada concluded that the risk of bladder cancer increased with both duration and concentration of DBP's in household water supplies (King and Marrett, 1996). In addition to the ingestion route, Villanueva et al. (2007) showed that inhalation and dermal absorption of THMs from household activities and swimming in pools may be associated with the development of cancer. A pooled analysis of six case-control studies, conducted by Villanueva et al. (1994), strengthened the conclusion that the risk of bladder cancer increases with long-term exposure to DBPs at levels that are currently observed in many

industrialized countries. In addition, a meta-analysis performed by Morris et al. (1992) concluded that there was a positive relationship between the consumption of DBP's in drinking water and bladder and rectal cancer in humans.

49. In recent years, U.S. EPA has set increasingly stringent limits on the allowable amounts of THMs in drinking water that determine what treatment facilities must achieve. In 1998, U.S. EPA issued the Stage 1 Disinfectants and Disinfection Byproducts Rule (Stage 1 DBPR), which reduced the maximum contaminant level (MCL) for THMs from 0.10 mg/L to 0.080 mg/L (100 to 80 parts per billion) and included additional compliance obligations for community water systems that chemically disinfect their water for primary or residual treatment (USEPA, 1998). Based upon results of testing following implementation of the Stage 1 rule, U.S. EPA subsequently determined that additional regulation beyond Stage I, in the form of the Stage 2 Disinfectant Byproducts Rule (Stage 2 DBPR), was needed to adequately protect human health (ADOH/DoE, 2003; USEPA, 2006c).

50. The Stage 2 DBPR places further restrictions on production and monitoring of DBPs, establishing revised Maximum Contaminant Levels (MCLs) and Maximum Contaminant Level Goals (MCLGs) for several of the DBPs. The concentration limits (MCLs) imposed by the Stage 2 DBP rule are, by definition, based upon a combination of considerations regarding health effects, technical feasibility and costs. In contrast, the development of MCLGs is based on health effects considerations only. For example, the MCLG for chloroform is 0.070 mg/L, while the MCL for total THMs, which typically are dominated by chloroform, is 0.080 mg/L. The MCLGs for two common HAA5s monochloroacetic acid and trichloroacetic acid, are 0.070 mg/L and 0.02 mg/L, respectively, while the current MCL for HAA5s is 0.060 mg/L. For the DBPs bromodichloromethane, bromoform, and dichloroacetic acid, the MCLG is zero, reflecting the ideal position that potential carcinogens should not be present in drinking water at any concentration. From a strictly health-based perspective (i.e., theoretical calculation using exposure guidelines) the following restrictive water concentrations were identified by USEPA (2006c) as being necessary to meet the standard regulatory benchmark of 1-in-one million cancer risk:

- dibromochloromethane 0.0009 mg/L
- bromodichloromethane 0.001 mg/L
- bromoform 0.008 mg/L
- dichloroacetic acid 0.0007 mg/L

In addition, the 2008 U.S. EPA Region 6 risk-based screening level for chloroform in residential water is 0.00017 mg/L (0.17 µg/L) (U.S. EPA, 2008). Wang (2007) has recently shown that water quality standards for THMs need to be reviewed to account for the different risks resulting from each individual THM species.

51. The State of Oklahoma requires that all public water systems monitor the finished water supplies for both total THMs (TTHM) and HAA5s. Available data through early-2008 from ODEQ was reviewed for the 18 currently active public water systems (Figure T2) that draw water from the Illinois River Watershed. There were 51 exceedances of the chloroform maximum contaminant level goal (MCLG) of 0.07 mg/L and 16 near exceedances (defined for purposes of this report as being within 10% of the MCLG). There were 56 exceedances of the TTHM standard of 0.08 mg/L and 29 near exceedances, and 35 exceedances of the HAA5s standard of 0.06 mg/L and 11 near exceedances. It should be noted that TTHM and HAA5s were not reported until late 2003. Prior to that time, when data for each individual component was available, TTHM values were estimated by summing these components. Currently, ODEQ has reported close to 30 violations for MCL average exceedances of either TTHM or HAA5s in tap water at customer facilities, or for sampling/reporting violations (ODEQ, 2008). There were also a multitude of exceedances of the risk-based criteria for bromodichloromethane, dibromochloromethane, and chloroform. Table T1 compares the exceedances between the risk based values and the listed MCL or MCLG values. Risk based values were exceeded over 90% of the time for both chloroform and bromodichloromethane, while the risk based value for dibromochloromethane was exceeded about 56% of the time. Stated simply, if risks of this magnitude were found at a waste disposal site or an industrial contamination site, in my experience they would require attention and remediation.

52. CDM collected water samples on three separate occasions in 2006 from 5 locations within three different IRW public water systems and analyzed them for TTHM and HAA5s. These locations are depicted in Figure T3. Of the total 45 samples collected, there were 6 exceedances and 5 near exceedances of the TTHM MCL (24%) and 3 exceedances and 1 near exceedance of the HAA5s MCL (9%). All of the samples exceeded the risk-based values for bromodichloromethane and dibromochloromethane and dichloroacetic acid, while two of the samples exceeded the individual chloroform MCLG of 0.070 mg/L. Six of the samples were greater than the MCLG of 0.02 mg/L for trichloroacetic acid. These results are illustrated in Table T2. In addition, raw water samples collected during the recreational season of 2005, 2006 and 2007 were evaluated for THM-forming potential (THMFP; CDM, 2008) and 71% of the results (57/80) showed values in excess of the TTHM MCL at twelve different locations along the Illinois River and in Lake Tenkiller. Table T3 shows the average THMFP from five PWAs. These report several of the larger systems which withdraw, treat and distribute water from the IRW.

53. Beyond the increased human health risks, elevated levels of THMs and HAA5s in drinking water often result in esthetic concerns (e.g., disagreeable taste and odors) in water supplies at concentrations which are at or near the drinking water standards (USEPA, 2006c). Thus, the water supply may be in compliance with regulatory numerical standards, but may not meet the Oklahoma narrative standards for water supplies. This general narrative criteria, found in OAC 785:45-5-9, states that “taste and odor producing substances from other than natural origin shall not interfere with the production of a potable water supply by modern treatment methods...” and “shall be maintained at all times to all surface waters of the state” (OAC, 2007). This criterion has not been met on many occasions in the IRW. In addition, the general criteria for public and private water supplies found in OAC 785:45-5-10 (5) (A) states that “The quality of the surface waters of the state which are designated as public or private water supplies shall be protected, maintained, and improved when feasible, so that the waters can be used as sources of public and private raw water supplies”. In this case the water bodies in the IRW are not being adequately protected. Also found in that criterion is OAC

785:45-5-10(5) (B) which states: “These waters shall be maintained so that they will not be toxic, carcinogenic, mutagenic, or teratogenic to humans” (OAC, 2007). The reported TTHM and HAA5s concentrations detected in IRW waters clearly demonstrate that this criterion is not being met.

54. The impacts from increased disinfection byproducts (e.g., THMs and HAA5s) associated with the application of poultry waste to lands in the Illinois River Watershed represent an imminent and substantial endangerment to human health.

Cyanobacteria

55. Cyanobacteria, also often termed “harmful blue-green algae”, are photosynthetic single-celled aquatic organisms that can produce potent toxins (e.g., microcystins, cylindrospermopsin, saxitoxins). It is well-documented that those toxins may be poisonous to aquatic organisms, to terrestrial animals, and to humans when present in water bodies at sufficient concentrations (Duy et al., 2000; Stewart, 2004; Falconer and Humpage, 2005; Janse et al., 2005; Health Canada, 2007; Cooke and Welch, 2008). The production of cyanobacterial toxins is influenced by a wide array of environmental conditions such as nutrient availability, light conditions, and temperature (Carson, 2000). Human health hazards occur from the following exposure routes: oral (through accidental ingestion of contaminated recreational water or ingestion of contaminated raw drinking water); dermal (through direct contact of exposed body parts with contaminated recreational waters); and inhalation (through airborne aerosols). Carmichael (2001) also reported exposure via the intravenous route in a dialysis clinic in Brazil as a result of contaminated local water supplies.

56. As ancient life forms often found living near the surface of warm water lakes and reservoirs, cyanobacteria are a well-recognized health hazard in many water supplies worldwide, including those in Europe, Australia, Asia and the Americas (WHO, 2000; Mankiewicz et al., 2005; Boaru et al., 2006; Stewart et al., 2006b). There are nearly 2,000 identified species of cyanobacteria, with more than 50 being recognized as potentially toxic to humans and animals (WHO, 2003). As noted in Chorus et al. (2000), “Cyanobacterial toxins occur naturally, but pollution with nutrients from agriculture

and domestic wastewater has led to increased fertilization (eutrophication) of many water bodies”.

57. The first reported cyanobacterial poisonings were documented as animal poisoning cases, while incidences of human toxicity have been reported more recently (WHO, 2003). The effects of cyanobacteria may include low level hay fever-like symptoms, progressing to skin rashes and gastrointestinal symptoms. In addition, the toxins released by cyanobacteria can cause more specific and severe hepatic and nervous system injury (Codd, 2000; Briand et al., 2003; Stewart et al., 2006a;). The occurrence of symptoms, as well as an elevated incidence of symptoms have been described with increasing duration of water contact, as well as with increasing cyanobacterial cell counts (Pilotto et al., 1997). This risk of adverse effects is related to exposure conditions, and it has been noted that “Swimmers involuntary swallow some water while swimming; in addition, bathing suits, particularly wet suits, could accumulate cyanobacterial material and enhance the disruption of cells and the liberation of cell content” (Mankiewicz et al., 2005). In water bodies that are governed by the PBCR requirements, such as those in the IRW, the State of Oklahoma mandates that such water bodies “shall not contain chemical, physical or biological substances in concentrations that are irritating to skin or sense organs or are toxic or cause illness upon ingestion by human beings” (OAC, 2007). Toxins produced by cyanobacteria fit that prohibition, and include anatoxins and saxitoxins (nervous system), as well as microcystins, nodularins and cylindrospermopsins, which are toxic to the liver (Carmichael et al., 2001). The chances of contracting illness from cyanobacteria have been reported to increase with increasing algal bloom density and often increase toward the end of a season when cells have reached greater density and then may encounter unfavorable environmental conditions, may “lyse” (burst), and release their toxins (Dittmann and Wiegand, 2006).

58. Cyanobacterial growth is known to be enhanced by increased nutrient loading with phosphorus and other nutrient enrichment from agriculture and wastewater to surface water bodies (QWQTF, 1992; WHO, 1999; Chorus et al., 2000; Kotak et al., 2000; WHO, 2000; Briand et al., 2003; Gobler et al., 2007; Health Canada, 2007; Cooke and

Welch, 2008). Stevenson (2008) found the nutrient concentration along the streams of the Illinois River Watershed to be high and to be directly related to poultry house density in the area. Many lakes and reservoirs worldwide develop high cyanobacterial cell concentrations, especially in the warm summer months, become demonstrably greenish in color, with algal species floating to the surface under warm, still conditions, forming scums with extreme cyanobacterial cell concentrations above one million cells per milliliter (1×10^6 cells/mL; Falconer and Humpage, 2005). Causes of nutrient input and fertilization, mainly from phosphorus, may include insufficiently treated sewage, agricultural runoff, manure application, effluent from intensive livestock industries and runoff from roads in urban areas (QWQTF, 1992; Chorus et al., 2000; Falconer and Humpage, 2005). Eutrophication, which is the biological response to the excess of nutrients into a water body, has been identified as a major water quality management issue in a number of countries (Codd, 2000; Cooke and Welch, 2008). In addition, when turbulence of the water column is low, cyanobacteria can build up dense populations based on temperature profiles, the nutrients present, and their ability to control cell buoyancy, thereby controlling their vertical position in the water column to optimize their growth conditions (Steinberg and Hartmann, 1988; Duy et al., 2000). As noted by Falconer and Humpage (2005): "There is a strong relationship between phosphorus concentration in the water and cyanobacterial numbers and also a similar though less linked relationship between dissolved nitrate/ammonia and cyanobacteria". Managing nutrient inputs to a water body has been shown to reduce the magnitude of algal biomass, as well as reducing the frequency and duration of algal proliferation events (Biggs, 2000). Although research is limited to this point, it has been suggested that bioaccumulation of cyanobacterial toxins in food supplies may lead to adverse impacts to human health and to natural ecosystems (Duy et al., 2000).

59. When cyanobacteria become dominant in the algal community of a water body, this exacerbates negative effects on water quality based upon factors such as reduced water clarity and transparency, high primary production of algae, decreased biodiversity, and potential oxygen depletion (Mankiewicz et al., 2005). Oxygen depletion [i.e., low dissolved oxygen (DO)] can cause massive fish kills and the release

of malodorous and unpleasant tasting compounds, as well as the release of the recognized cyanobacterial toxins. Surface aggregations of cyanobacteria are possible due to their capacity to regulate their buoyancy, enabling them to localize preferentially at water depths where conditions are optimal for their growth (WHO, 2000; Dittmann and Wiegand, 2006). Visible scums of algae have been reported on the surface of Lake Tenkiller and historical data show that cyanobacteria often are the dominant phytoplankton community in the reservoir (Cooke and Welch, 2008).

60. Aside from producing toxins, cyanobacteria can be a visual nuisance and blooms often reduce water clarity, resulting in decreased recreational value of impacted lakes and reservoirs (Chorus et al., 2000; Kotak and Zurawell, 2007). The presence of cyanobacteria also can cause taste/odor problems from a number of chemicals they release, such as geosmin and methylisoborneol (Watson et al., 2003; Tung et al., 2004; Dittmann and Wiegand, 2005; Izaquirre and Taylor, 2007; Uwins et al., 2007; Juttner, 2007; Xie et al., 2007; Cooke and Welch, 2008). In addition to being free of toxic materials, according to the Oklahoma Administrative Code (OAC) 785:45-5-19 (regarding surface water supplies): “The water must also be free from noxious odors and tastes, from materials that settle to form objectional deposits, and discharges that produce undesirable effects or are a nuisance to aquatic life” (OAC, 2007). Results from a telephone survey conducted in 2006 on behalf of the state of Oklahoma, among personnel from 20 utilities along the Illinois River and Lake Tenkiller, indicated that five of the utilities (Burnt Cabin, Cherokee Co. RWD #2, Cherokee Co. RWD #13, Pettit Mountain Water, and Tenkiller Aqua Park) had received complaints of taste and odor, thus violating this Oklahoma standard. These taste and odor complaints occurred during the summer and fall months. Increases in the utilities’ usage of alum, activated carbon and/or chlorine reportedly were employed to deal with this problem. Utilities with the most complaints are located in the upper reservoir, where eutrophic to hypereutrophic conditions have been shown to occur during the summer months (Cooke and Welch, 2008). Potential explanations for these negative taste and odor complaints include cyanobacterial algal products.

61. The measurement of “chlorophyll a” (Chl) concentration is used as another indicator of the abundance of algal growth (including cyanobacteria) in water bodies. Chl concentrations are used to define trophic state (or nutrient content) of water bodies and can be related to taste and odor issues. Cooke and Welch (2008) noted that the probability of cyanobacteria “blooms” rise sharply when mean summer Chl concentration is above 10 µg/L. Two sources of lake water odors that reasonably may be related to trophic state (e.g., eutrophication) have been described as high concentrations of specific algal species and bottom water odors that result from substances released from the sediment under anoxic (i.e., low oxygen) conditions (Arruda and Fromm, 1989). Existing regulations [OAC 785:45-5-10 (7)] describe the numerical criterion for Chl as “The long term average concentration of chlorophyll-a at a depth of 0.5 meters below the surface shall not exceed 0.010 milligrams per liter” in Lake Tenkiller (OAC, 2007). In this instance, “long-term average” is defined as the “arithmetic mean of at least 10 samples taken across at least 12 months” (OAC, 2007). Results from CDM sampling, as shown in Table C1, at four locations on Lake Tenkiller show that this long-term average criterion was exceeded during 2005-2006 at two of the four locations sampled with the long-term averages of 15.9 and 27.1 µg/L at LK-03 and LK-04, respectively (CDM, 2008). The other two locations, LK-01 and LK-02, had long-term average values of 6.4 and 8.7 µg/L, respectively. Although there were not enough samples to calculate long-term averages from the acquired Oklahoma Water Resources Board (OWRB) Chl data, calculated averages from 1999 through 2006 (between 10 and 15 samples at each site) at 7 locations on Lake Tenkiller show levels above 10 µg/L Chl at four of the locations (OWRB, 2006). These seven locations are illustrated in Figure C1.

62. WHO (1999) guidelines are widely used in the public health community to evaluate potential risks that may be posed by cyanobacteria in water supplies. Those guidelines state that at a density less than 20,000 cells/ mL of cyanobacteria there exists relatively low risk of adverse human health effects, while at or above 100,000 cells/mL there is moderate risk and, when visible scum of cyanobacteria is present, there is high risk of adverse health effects. Pilotto (1997) has suggested that these guidelines, while

they are commonly employed, may not in and of themselves be adequate to protect human health.

63. Results from sampling by CDM (2006, 2007) and OWRB and U.S. Army Corps of Engineers (ACoE; 2004, 2005) from several different locations (Figure C1 and C2) on Lake Tenkiller during August 2004 through August 2007 showed that approximately 58% of all samples (233/404) exhibited cyanobacterial densities of greater than 20,000 cells/mL. In addition, as listed in Table C2, approximately 24% (55/233) of those samples that were greater than 20,000 cells/ml exceeded 100,000 cells/mL (moderate risk), and one June 2006 sample exceeded 1,000,000 cells/mL.

64. Cell sizes of the cyanobacteria can vary considerably within and between species, therefore cell numbers alone may not be the only or the ideal measure of population size or potential toxicity risk (WHO, 1999). It is possible for biological volume of cells ("biovolume") to be estimated from cell counts and average cell volumes, and the resultant converted data reported as micrometers cubed per milliliter of water ($\mu\text{m}^3/\text{mL}$). However, no federal or state standards currently are available for comparison with measured biovolume values. When compared to measuring cyanobacterial genotypes, Janse et al. (2005) found that the monitoring of algal biomass/biovolume can be an adequate predictor of toxin production. The ACoE collected samples during the period from 2001 through 2004 from four different locations along Lake Tenkiller (Figure C2), and reported cyanobacteria in units of biovolume. These results are shown in Table C3 and show biovolume levels ranging from 1,800 to 60,802,075,714 $\mu\text{m}^3/\text{mL}$ (over 60 billion $\mu\text{m}^3/\text{mL}$). Of the 128 total samples, 64% were greater than 1,000,000 $\mu\text{m}^3/\text{mL}$, 28% were greater than 100,000,000 $\mu\text{m}^3/\text{mL}$, and 20% were greater than 1,000,000,000 $\mu\text{m}^3/\text{mL}$. Cyanobacteria biovolume ranges for all of the CDM and OWRB samples that were greater than 20,000 cells/mL ranged from 22,000 $\mu\text{m}^3/\text{mL}$ to 13,000,000 $\mu\text{m}^3/\text{mL}$. Thus, despite the absence at present of a rigorous biovolume standard, these results clearly represent potentially harmful amounts of cyanobacterial algal growth along Lake Tenkiller over four consecutive seasons.

65. The demonstrated increases in algal growth in parts of the Illinois River Watershed and Lake Tenkiller represent an imminent and substantial endangerment to human health.

Statement of Opinions

66. Based on the information presented in this report, coupled with my professional training and experience, I have the following opinions:

- There are ongoing and widespread conditions of bacterial contamination elevated over a large proportion of the Illinois River Watershed, including those areas where common uses of the Illinois River and its tributaries include aquatic recreation. The magnitude and the distribution of the elevated bacterial contamination represent imminent and substantial endangerments to human health. Several lines of evidence support the conclusion that these elevated levels of bacteria are primarily attributable to the land disposal of poultry waste.
- Increases in nutrients (e.g., phosphorus) related to the land disposal of poultry waste have resulted in eutrophication and increased algal growth broadly in the Illinois River Watershed, including Lake Tenkiller. These increased levels of algae and other forms of waterborne organic carbon combine with the normal drinking water disinfection process to produce potentially dangerous Disinfection Byproducts, such as trihalomethanes (THMs) and haloacetic acids (HAA5s). Routine and specific sampling results have identified levels of THMs and HAA5s in drinking water distribution systems that withdraw water from the Illinois River Watershed, and these levels represent an imminent and substantial endangerment to human health.
- Increases in nutrients (e.g., phosphorus) related to the land disposal of poultry waste have resulted in eutrophication and increased algal growth broadly in the Illinois River Watershed, including Lake Tenkiller. Among the measures of increased eutrophication is the detection of potentially dangerous levels of Cyanobacteria, also termed "Harmful Blue Green Algae". The levels of Cyanobacteria reported in a number of studies conducted within Lake Tenkiller represent an imminent and substantial endangerment to human health.
- There are biological and chemical hazards and impairments (i.e., bacteria and indicator organisms, THMs/HAA5s, cyanobacteria) within the Illinois River Watershed which are present at levels that are capable of causing damage to human health and which will continue to do so unless action is taken to eliminate the sources or major contributing factors for each of these hazards and impairments. The process of applying poultry waste to fields is a significant contributor to the development and persistence of

these hazards and impairments of the IRW. The hazards and impairments represent an imminent and substantial endangerment to human health.


67. I reserve the right to modify and supplement the opinions presented in this report, to the extent that I become aware of new information, if reinterpretation of existing information occurs, or if it is appropriate to provide response and/or rebuttal to the technical opinions of other experts in this matter.


68. Compensation for all of my professional activities in this matter is at a rate of \$195 per hour.

69. A list of the technical references that are cited herein is included as Attachment A to this report.

70. A complete copy of my Curriculum Vitae is included as Attachment B to this report.

71. A list of sworn testimony that I have provided in deposition or at trial for the period from May 15, 2004 through May 15, 2008 is presented as Attachment C to this report.


CHRISTOPHER M. TEAF, Ph.D.
Toxicologist and Risk Assessment Specialist


DATE

ATTACHMENT A

References Cited

REFERENCES CITED

- Adamski, J.C. 1987. The effect of agriculture on the quality of ground water in karstified carbonate terrain, Northwest Arkansas. Master of Science thesis for the University of Arkansas.
- Adamski J.C. and Steele, K.F. 1988. Agricultural land use effects on ground water quality in the Ozark region. Proceedings of Agricultural Impacts on Groundwater Conference. Dublin, OH: National Water Well Association. p. 593-614.
- Adegoke, G.O. 1986. Characteristics of staphylococci isolated from man, poultry and some other animals. *J Appl Bacteriology*, 60:97-102.
- Allos, B.M. 2001. *Campylobacter jejuni* infections: Update on emerging issues and trends. *Clin Infect Dis* 34(5):719-720..
- Anderson, D.M. et al. 2002. Harmful algal blooms and eutrophication: Nutrient sources, composition, and consequences. *Estuaries* 25(4b):704-726.
- Arkansas Department of Health, Division of Engineering (ADOH/DoE). 2003. Arkansas Drinking Water Update. Volume 16, Number 3.
- Arruda, J.A. and C.H. Fromm. 1989. The relationship between taste and odor problems and lake enrichment from Kansas lakes in agricultural watersheds. *Lake and Reservoir Management* 5(1):45-52.
- Barbour, E.K. and N.H. Nabbut. 1981. Isolation of Salmonella and some other potential pathogens from two chicken breeding farms in Saudi Arabia. *Avian Diseases* 26(2):234-244.
- Barrell, R.A.E. et al. 2000. Microbiological standards for water and their relationship to health risk. *Communicable Disease and Public Health* 3(1):8-13.
- Bass, L. et al. 1999. Incidence and characterization of integrons, genetic elements mediating multiple-drug resistance, in avian *Escherichia coli*. *Antimicrobial Agents and Chemo* 43(12): 2925-2929.
- Belanger, A. and T. Shryock. 2007. Macrolide-resistant *Campylobacter*: the meat of the matter. *J Antimicrobial Chemotherapy* 60:715-723.
- Biggs, B.J.F. 2000. Eutrophication of streams and rivers: dissolved nutrient-chlorophyll relationships for benthic algae. *J N Am Benthol Soc* 19(1):17-31.
- Blackburn, B.G. et al. 2004. Surveillance for waterborne-disease outbreaks associated with drinking water- United States, 2001-2002. *MMWR* 53(No. SS-8):23-45.
- Boaru, D.A. et al. 2006. Toxic potential of microcystin-containing cyanobacteria extracts from three Romanian freshwaters. *Toxicon* 47:925-932.
- Brabazon E.D. et al., 2007. Under-reporting of notifiable infectious disease hospitalizations in a health board region in Ireland: room for improvement? *Epidemiol Infect* 136:241-247.
- Bren, L. 2001. Antibiotic Resistance from down on the chicken farm. USDA/FDA Consumer Magazine, Jan-Feb. 2001.

- Briand, J.F. et al. 2003. Health hazards for terrestrial vertebrates from toxic cyanobacteria in surface water ecosystems. *Vet Res* 34: 361-377.
- Brosnan, T.M. et al. 1996. Long term improvements in the water quality due to sewage abatement in the Lower Hudson River estuaries. *Estuaries* 19(4): 890-900.
- Brown, L.R. 2003. Potential effects of organic carbon production on ecosystems and drinking water quality. San Francisco Estuary and Watershed Science, 1(1), Paper #3:1-22.
- Bull, R.J. et al. 1995. Water Chlorination: Essential process or cancer hazard? *Fund Appl Toxicol* 28:155-166.
- Byappanahalli, M.N. et al. 2003. Growth and survival of *Escherichia coli* and enterococci populations in the macro-alga *Cladophora* (Chlorophyta). *FEMS Microbiology Ecology* 46:203-211.
- Camp, Dresser & McKee (CDM). 2008. Personal communications with technical staff and data tables sent from staff over the length of the project.
- Caneday, L. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Cantor, K.P. et al. 1978. Associations of cancer mortality with halomethanes in drinking water. *J Natl Cancer Inst* 61(4):979-985.
- Carmichael, W. et al. 2001. Human fatalities from cyanobacteria: Chemical and biological evidence for cyanotoxins. *Environ Health Perspect* 109:663-668.
- Carson, B. 2000. Cyindrospermopsin- a review of toxicological literature. Prepared by Integrated Laboratory Systems for Scott Masten, National Institute of Health Sciences, Research Triangle Park, NC. Contract number N01-ES-65402.
- Centers for Disease Control and Prevention (CDC). 2008. *E. coli*. <http://www.cdc.gov/ecoli/>.
- Chen, L. et al. 2006. Dose Response for infectivity of several strains of *Campylobacter jejuni* in chickens. *Risk Analysis* 26(6):1613-1621.
- Chorus, I. et al. 2000. Health risks caused by freshwater cyanobacteria in recreational waters. *J Toxicol Environ Health* 3(4):323-347.
- Codd, G.A. 2000. Cyanobacterial toxins, the perception of water quality, and the prioritization of eutrophication control. *Ecol Eng* 16:51-60.
- Cooke, G.D. and E.B. Welch. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Coyne, M.S. and R.L. Blevins. 1995. Fecal bacteria in surface runoff from poultry-manured fields. *Animal Waste and the Land-Water Interface*. Steele, K. (Ed.). Boca Raton, FL: Lewis Publishers.
- Crane, S.R. et al. 1980. Die-off of fecal indicator organisms following land application of poultry manure. *J Environ Qual* 9(3):531-537.

- Craun, G.F. and R.L. Calderon. 2006. Workshop summary: estimating waterborne disease risks in the United States. *J Water and Health* 4(2):241-253.
- Davis, R. et al. 2005. *Escherichia Coli* survival in Mantled Karst Springs and streams, northwest Arkansas Ozarks, USA. *J. Am. Water Resources Assn*, Dec. 2005: 1279-1287.
- Diarra, M.S. et al. 2007. Impact of feed supplementation with antimicrobial agents on growth performance of broiler chickens, *Clostridium perfringens* and *Enterococcus* counts, and antibiotic resistance phenotypes and distribution of antimicrobial resistance determinants in *Escherichia coli* isolates. *Appl Environ Microbiol* 73(20):6566-6576.
- Dipineto, L. et al. 2006. Presence of Shiga toxin-producing *Escherichia coli* 0157:H7 in living layer hens. *Letters in Appl Microbiol* 43:293-295.
- Dittmann, E. and C. Wiegand. 2006. Cyanobacterial toxins- occurrence, biosynthesis and impact on human affairs. *Mol Nutr Food Res* 50:7-17.
- Doane, C.A. et al. 2007. Occurrence of *Escherichia coli* 0157:H7 in diverse farm environments. *J Food Prot* 70(1):6-10.
- Dore, M. et al. 1982. Reactivity of halogens with aqueous micropollutants: A mechanism for the formation of trihalomethanes. *J Am Water Works Assn* 74(2):103-107.
- Doyle, M.P. and M.C. Erickson. 2006. Reducing the carriage of foodborne pathogens in livestock and poultry. *Poultry Science* 85:960-973.
- Doyle, M.P. and J.L. Schoeni. 1987. Isolation of *Escherichia coli* 0157:H7 from retail fresh meats and poultry. *Appl Environ Microbiol* 53(10):2394-2396.
- Dupont, H.L. 2007. The growing threat of foodborne bacterial enteropathogens of animal origin. *Food Safety* 45:1353-61.
- Duy, N. et al. 2000. Toxicology and risk assessment of freshwater cyanobacteria (blue-green algal) toxins in water. *Rev Environ Contam Toxicol* 163:113-186.
- Ellis-Pegler, R.B. et al. 1995. A placebo controlled evaluation of Lomefloxacin in the treatment of bacteria diarrhea in the community. *J Antimicrobial Chemotherapy* 36(1):259-263.
- Engel, B. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Falconer, I.R. and A.R. Humpage. 2005. Health risk assessment of cyanobacterial (blue-green algal) toxins in drinking water. *Intl J Environ Res Public Health* 2(1):43-50.
- Fisher, J. B. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Fruh, E.G. 1967. The overall picture of eutrophication. *J WPCF* 39:1449-1463.
- Gagliardi, J.V. and J.S. Karns. 2000. Leaching of *Escherichia coli* O157:H7 in diverse soils under various agricultural management practices. *Appl Environ Microbiol* 66(3):877-863.

- Gallard, H. and U. Gunten. Chlorination of natural organic matter: kinetics of chlorination and of THM formation. *Water Res* 36:65-74.
- Giddens, J. and A.P. Barnett. 1980. Soil loss and microbiological quality of runoff from land treated with poultry litter. *J Environmental Quality* 9(3):518-520.
- Gobler, C.J. et al. 2007. The contrasting impacts of nutrient loading and zooplankton grazing on the growth and toxicity of cyanobacteria blooms in a eutrophic New York lake. www.epa.gov/cyano_habs_symposium/abstract.
- Gopal, K. et al. 2007. Chlorination byproducts, their toxicodynamics and removal from drinking water. *J Hazardous Materials* 140:1-6.
- Hald, B. et al. 2007. Use of fly screens to reduce *Campylobacter* spp. Introduction in boiler houses. *Emerging Infectious Disease* 13(12):1951-1953.
- Hartel, P.G. et al. 2000. Survival of fecal coliforms in fresh and stacked broiler litter. *J Appl Poultry Res* 9:505-512.
- Harwood, V.J. et al. 2005. Validity of the indicator organism paradigm for pathogen reduction in reclaimed water and public health protection. *Appl Environ Microbiol* 71(6):3163-3170.
- Harwood, V. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Health Canada. 2007. Blue-green algae (cyanobacteria) and their toxins. http://www.hc-sc.gc.ca/ewh-semt/water-eau/drink-potab/cyanobacteria-cyanobacteries_e.html.
- Himathongkham, S. et al. 1999. Survival of *Salmonella enteritidis* and *Salmonella typhimurium* in chicken manure at different levels of water activity. *FEMS Microbiol Letters* 172:159-163.
- Hoehn, R.C. et al. 1980. Algae as sources of trihalomethane precursors. *J Am Water Works Assn* 72(6):344-350.
- Hofacre, C.L. et al. 1999. Presence of fluoroquinolone-resistant coliforms in poultry litter. *Avian Diseases* 44:963-967.
- Hurd, H.S. et al. 2004. Public health consequences of macrolide use in food animals: a deterministic risk assessment. *J Food Prot* 67(5): 980-992.
- Ishii, S. et al. 2006. *Cladophora* (Chlorophyta) spp. harbor human bacterial pathogens in nearshore water of Lake Michigan. *Appl Environ Microbiol* 72(7):4545-4553.
- Izaguirre, G. and W.D. Taylor. 2007. Geosmin and MIB events in a new reservoir in southern California. *Water Science and Technology* 55(5):9-14.
- Jacobs-Reitsma, W.F. et al. 1994. Cecal carriage of *Campylobacter* and *Salmonella* in Dutch broiler flocks at slaughter: a one-year study. *Poultry Science* 73:1260-1266.
- Jamieson, R.C. et al. 2002. Movement and persistence of fecal bacteria in agricultural soils and subsurface drainage water: A review. *Canadian Biosystems Engineering* 44:1.1-1.9.

- Janse, I. et al. 2005. Contrasting microcystin production and cyanobacterial population dynamics in two Planktothrix-dominated freshwater lakes. *Environ Microbiol* 7(10):1514-1524.
- Jenkins, M.B. et al. 2006. Fecal bacteria and sex hormones in soil and runoff from cropped watersheds amended with poultry litter. *Sci Total Environ* 358:164-177.
- Juttner, F. 2007. Needs and perspectives of odour research in the aquatic sciences. *Water Science and Technology* 55(5):367-369.
- Kazwala, R.R. et al. 1990. Factors responsible for the introduction and spread of *Campylobacter jejuni* infection in commercial poultry production. *Vet Record* 126:305-306.
- Kelley, T.R. et al. 1995. Bacterial pathogens and indicators in poultry litter during re-utilization. *J Appl Poultry Res* 4:366-373.
- King W.D. and L.D. Marrett. 1996. Case-control study of bladder cancer and chlorination by-products in treated water (Ontario, Canada). *Cancer Causes and Controls* 7(6):596-604.
- Kloos, W.E. and T.L. Bannerman. 1999. *Staphylococcus and Micrococcus*. Manual of Clinical Microbiology, 7th edn. ASM Press. Washington, D.C.
- Kotak, B. et al. 2000. Role of chemical and physical variables in regulating microcystin-LR concentration in phytoplankton of eutrophic lakes. *Canad J Fish Aquatic Sci* 57:1584-1593.
- Kotak, B. and R. Zurawell. 2007. Cyanobacterial toxins in Canadian freshwaters: A review. *Lake and Reservoir Management* 23:109-122.
- Krasner, S.W. et al. 2006. Occurrence of a new generation of disinfection byproducts. *Environ Sci Technol* 40:7175-7185.
- Leclerc, H. et al. 2002. Microbial agents associated with waterborne disease. *Critical Reviews in Microbiology* 28(4):371-409.
- Lee, G. F. and R.A. Jones. "Regulating Drinking Water Quality at the Source," Proc. University of California Water Resources Center Conference: Protecting Water Supply Water Quality at the Source, Sacramento, CA, 39pp, April (1991).
- Lee, S.H. et al. 2002. Surveillance for waterborne-disease outbreaks— United States, 1999-2000. *MMWR*, 2002: 51(No. SS-8):1-28.
- Lee, W. et al. 2007. Dissolved organic nitrogen as a precursor for chloroform, dichloroacetonitrile, N-nitrosodimethylamine, and trichloronitromethane. *Environ Sci Technol* 41:5485-5490.
- Li, X. et al. 2006. *Salmonella* populations and prevalence in layer feces from commercial high-rise houses and characterization of the *Salmonella* isolates by serotyping, antibiotic resistance analysis, and pulsed field gel electrophoresis. *Poultry Science* 86:591-597.
- Liang, J.L. et al. 2006. Surveillance for waterborne-disease and outbreaks associated with drinking water and water not intended for drinking- United States, 2003-2004. *MMWR*, 2006: 55(No. SS-12):31-58.

- Lindblom, G.B. et al. 1986. Natural *Campylobacter* colonization in chickens raised under different environmental conditions. *J of Hygiene* 96(3):385-391.
- Mankiewicz, J. et al. 2005. Hepatotoxic cyanobacterial blooms in the lakes of Northern Poland. *Environ Toxicol* 20:499-506.
- Marhaba, T.F. and I.H. Kochar. 2000. Rapid prediction of disinfection by-product formation potential by fluorescence. *Environ Eng Policy* 2:29-36.
- Martin, A.B. et al. 1993. Lake sediments as potential sources of trihalomethane precursors. *Water Res* 27(12):1725-1729.
- Mayo Clinic. 2008. *Salmonella*. <http://www.mayoclinic.com/health/salmonella/DS00926>.
- Metcalf & Eddy. 1991. Wastewater Engineering: Treatment, Disposal and Reuse. New York. McGraw-Hill Publishing. pp: 108-111.
- Morris, R.D. et al. 1992. Chlorination, chlorination by-products, and cancer: A meta-analysis. *Am J Public Health* 82(7):955-963.
- Murray, P.R. et al. 2003. Manual of Clinical Microbiology (Eighth Edition). Washington, DC: ASM Press.
- Nandi, S. et al. 2004. Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistant integrons in poultry litter. *Proc Natl Acad Sci* 101(18): 7118-7122.
- National Academy of Sciences (NAS). 1987. Drinking Water and health (Vol. 7): Disinfectants and Disinfectant By-Products. Washington, DC: National Academy Press.
- National Oceanic and Atmospheric Administration (NOAA). 2000. Climatography of the United States. Contained in <http://cdo.ncdc.noaa.gov/climatenormal/clim81/oknorm.pdf>.
- National Research Council (NRC). 2004. Indicators for waterborne Pathogens. National Research Council for the National Academies. National Academies Press. Washington, DC.
- Neill, S.D. et al. 1984. *Campylobacter* species in broiler chickens. *Avian Pathology* 13:777-785.
- Nokes, C.J. et al. 1999. Modeling the formation of brominated trihalomethanes in chlorinated drinking waters. *Wat Res* 33(17):3557-3568.
- Office of the Secretary of Environment (OSE). 2003. Coordinated watershed restoration and protection strategy for Oklahoma's impaired scenic rivers. January 31, 2003.
- Oklahoma Administrative Code (OAC). 2007. Title 785. Oklahoma Water Resources Board, Chapter 45. Oklahoma's water quality standards.
- Oklahoma Conservation Commission (OCC). 2004. Illinois River and Baron Fork Watershed Implementation Project. OCC Task 113. FY 1999 319(h) Task 800. Final Report. December 2004.
- Oklahoma Department of Agriculture (ODAFF). 2008. Agriculture Laws and Statutes webpage. <http://www.oda.state.ok.us/ogc-laws.htm>.

Oklahoma Department of Environmental Quality (ODEQ). No date. Oklahoma Total Maximum Daily Load Practitioners Guide.

Oklahoma Department of Environmental Quality (ODEQ). 2006a. The State of Oklahoma Water Quality Assessment Integrated Report.

Oklahoma Department of Environmental Quality (ODEQ). 2008. ODEQ Public Water Supply website.

Oklahoma State Department of Health (OSDOH), Communicable Disease Division. 2006. Personal communications with data management staff.

Oklahoma State Department of Health (OSDOH). 2007. Information presented with associated background at <http://www.health.state.ok.us>

Oklahoma Water Resources Board (OWRB). Greater Tenkiller Area Association and Illinois River Oklahoma Water Watch Chapters: Draft 1997-2002 Data Summary.

Oklahoma Water Resources Board (OWRB). 2005. 2005 Report of the Oklahoma beneficial use monitoring program streams report.

Oklahoma Water Resources Board (OWRB). 2007. Data sets for samples taken from the Illinois River. Unpublished.

Oklahoma Water Resources Board (OWRB). 2008. Reported well log viewer database at <http://www.owrb.ok.gov/maps/server/wims.php>.

Oliver, B.G. and J. Lawrence. 1979. Haloforms in drinking water: A study of precursors and precursor removal. *J Am Water Works Assoc* 71:161-163.

Olsen, R. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.

O'Reilly, C.E. et al. 2007. A Waterborne outbreak of gastroenteritis with multiple etiologies among Resort Island visitors and residents: Ohio, 2004. *CID* 44:506-512.

Peterson, H.G. et al. 1993. Trihalomethanes in finished drinking water in relation to dissolved organic carbon and treatment process for Alberta surface waters. *Environ Tech* 14:877-884.

Pew Commission on Industrial Farm Animal Production (PCIFAP). 2008. Putting meat on the table: Industrial farm animal production in America. A project of The Pew Charitable Trusts and Johns Hopkins Bloomberg School of Public Health.

Pilotto, L.S. et al. 1997. Health effects of exposure to cyanobacteria (blue-green algae) during recreational water-related activities. *Australian and New Zealand Journal of Public Health* 21(6):562-566.

Preusch, P.L. et al. 2002. Nitrogen and phosphorus availability in composted and uncomposted poultry litter. *J Environ Qual* 31:2051-2057.

Price, L.B. et al. 2007. Elevated risk of carrying gentamicin-resistant *Escherichia coli* among U.S. poultry workers. *Environ Health Perspect* 115(12):1738-1742.

- Pruss, A. 1998. Review of epidemiology studies on health effects from exposure to recreational water. *International J Epidemiology* 27:1-9.
- Queensland Water Quality Task Force (QWQTF). 1992. Report on freshwater algal blooms in Queensland. Vol 1 Main Report. Queensland Water Quality Task Force. Queensland Government.
- Reckhow, D.A. and P.C. Singer. 1990. Chlorination by-products in drinking waters: from formation potentials to finished water concentrations. *J Am Water Works Assoc* 82:173-180.
- Richardson, S.D. et al. 2007. Occurrence, genotoxicity, and carcinogenicity of regulated and emerging disinfection by-products in drinking water: a review and roadmap for research. *Mutat Res* 636(1-3):178-242.
- Ringbauer, J. et al. 2006. Effects of large-scale poultry farms on aquatic microbial communities: A molecular investigation. *J Water Health* 41(1):77-86.
- Rook, J.J. 1976. Haloforms in drinking water. *J AWWA* 68:168-172.
- Sas, H. (Ed.). 1989. Lake restoration by reduction of nutrient loading: Expectations, experiences, extrapolations. Academia Verlag Richarz, St. Augustin, Germany. p. 497.
- Slaton, N.A. 2004. Nutrient input and removal trends for agricultural soils in nine geographic regions in Arkansas. *J Environ Qual* 33:1606-1615.
- Smith, J.L. et al. 2007. Impact of usage on antimicrobial resistance in commercial *E. coli* strains colonizing broiler chickens. *Appl Environ Microbiol* 73(5):1404-1414.
- Steinberg, C.E.W. and H.M. Hartmann. 1988. Planktonic bloom-forming Cyanobacteria and the eutrophication of lakes and rivers. *Freshwater Biology* 20(2):279-287.
- Stepczuk, C. et al. 1998. Allochthonous contributions of THM precursors to a eutrophic reservoir. *Journal of Lake and Reservoir Management* 14(2-3):344-355.
- Stevens, A. et al. 1976. Chlorination of organics in drinking water. *J AWWA* 68(11):615-620.
- Stevenson, R.J. 2008. Draft Report. State of Oklahoma vs. Tyson Foods, et al. United States District Court, Northern District of Oklahoma. Case Number: 05-CV-329-GKF-SAJ.
- Stewart, I. 2004. Recreational exposure to freshwater cyanobacteria: epidemiology, dermal toxicity, and biological activity of cyanobacterial lipopolysaccharides. Dissertation, University of Queensland, Australia.
- Stewart, I. et al. 2006a. Cutaneous hypersensitivity reactions to freshwater cyanobacteria – human volunteer studies. *Dermatology* 6(1): 6-16.
- Stewart I. et al. 2006b. Recreational and occupational field exposure to freshwater cyanobacteria- a review of anecdotal and case reports, epidemiological studies and the challenges for epidemiologic assessment. *Environmental Health: A Global Access Science Source*, 5:6.

- Stuart, M.E. et al. 1999. Trihalomethane formation potential: A tool for detecting groundwater contamination and the health risks associated with chlorination. Proceed. Sardinia 99, Seventh International Waste Management Landfill Symposium.
- Tang, Z. et al. 2005. Forecasting land use change and its environmental impact at a watershed scale. *J Environ Manage* 76:35-45.
- Terzich, M. et al. 2000. Survey of pathogens in poultry litter in the United States. *J Appl Poultry Res* 9:287-291.
- Tetra Tech. 2004. Manure Management. EPA Regional Priority AFO Science Question Synthesis Document. College Park, Maryland.
- Toranzos, G.A. et al. 2002. Detection of indicator microorganisms in environmental freshwaters and drinking waters. In: Manual of Environmental Microbiology. Hurst, C. (Ed.). Washington D.C.: ASM Press.
- Trampel, D.W. et al. 2007. Characterization of *Escherichia coli* isolates from peritonitis lesions in commercial laying hens. *Avian Diseases* 51:840-844.
- Tung, S.C. et al. 2004. The effect of oxidants on 2-MIB concentration with the presence of cyanobacteria. *Water Science and Technology* 49(9):281-288.
- University of Missouri Vet Med Diagnostic Lab. 2007. Avian necropsy. September 12, 2007. Unpublished. Bates #: CARTP166801 through CARTP166807.
- U.S. Census Bureau. 1990. Population Census database.
- U.S. Census Bureau. 2000. Population Census database.
- United States Department of Agriculture (USDA). 1991. Campylobacter questions and answers. USDA Food Safety and Inspection Service background Document, May 1991.
- U.S. Environmental Protection Agency (USEPA). 1984. Health Effects Criteria for Fresh Recreational Waters. EPA-600/1-84-004.
- U.S. Environmental Protection Agency (USEPA). 1986. Ambient Water Quality Criteria for Bacteria. EPA440/5-84-002.
- U.S. Environmental Protection Agency (USEPA). 1997. Compendium of Tools for Watershed Assessment and TMDL Development. EPA 841-B-97-006. U.S. Environmental Protection Agency. Office of Wetlands, Oceans, and Watersheds, Washington, D.C.
- U.S. Environmental Protection Agency (USEPA). 1998. Stage 1 disinfectants and disinfection by-products rule. EPA 815-F-98-010.
- U.S. Environmental Protection Agency (USEPA). 2001a. National Land Cover Data (NLCD) database.
- U.S. Environmental Protection Agency (USEPA). 2001b. Protocol for developing pathogen TMDLs. EPA 841-R-00-002. U.S. Environmental Protection Agency. Office of Water, Washington, D.C.

- U.S. Environmental Protection Agency (USEPA). 2003. Draft Implementation Guidance for Ambient Water Quality Criteria for Bacteria.
- U.S. Environmental Protection Agency (USEPA). 2005. Detecting and Mitigating the Environmental Impact of Fecal Pathogens Originating from Confined Animal Feeding Operations: Review. EPA/600/R-06/021.
- U.S. Environmental Protection Agency (USEPA). 2006a. National Primary Drinking Water Regulations: Groundwater Rule.
- U.S. Environmental Protection Agency (USEPA). 2006b. National Primary Drinking Water Regulations: Long Term 2 Enhanced Surface Water Treatment Rule: Final Rule.
- U.S. Environmental Protection Agency (USEPA). 2006c. Stage 2 disinfectants and disinfection by-products rule. Federal Register 71(2):388-492.
- U.S. Environmental Protection Agency (USEPA). 2007a. Integrated Risk Information System (IRIS) database.
- U.S. Environmental Protection Agency (USEPA). 2007b. Permit Compliance System (PCS) database.
- U.S. Environmental Protection Agency (USEPA). 2007c. Report of the experts scientific workshop on critical research needs for the development of new or revised recreational water quality criteria. EPA 823-R-07-006. U.S. Environmental Protection Agency. Office of Water, Office of Research and Development, Warrenton, Virginia.
- U.S. Environmental Protection Agency (USEPA). 2008. Region 6 Human Health Medium-specific Screening Levels.
- U.S. Food and Drug Administration Ctr. for Drug Eval. & Research (USFDA). 2002. Hearing for the anti-infective drugs advisory committee. February 20, 2002.
- U.S. Geological Service (USGS). 2007a. Data sets for samples taken from the Illinois River. Unpublished.
- U.S. Geological Service (USGS). 2007b. Bacteria and their effects on ground-water quality. <http://mi.water.usgs.gov/h2oqual/GWBactHOWeb.html>.
- Uwins, H.K. et al. 2007. A case study investigating the occurrence of geosmin and 2-methylisoborneol (MIB) in the surface waters of the Hinze Dam, Gold Coast, Australia. *Water Science and Technology* 55(5):231-238.
- Van Steenderen, R.A. et al. 1988. Microcystis aeruginosa as an organohalogen precursor. *Water SA* 14(1):59-62.
- Victorian Government Health Information (VGHI). 2007. Food or water-borne illness. Contained in http://www.health.vic.gov.au/ideas/bluebook/food_water.
- Villanueva, C.M. et al. 1994. Disinfection byproducts and bladder cancer. *Epidemiology* 15(3):357-367.
- Villanueva, C.M. et al. 2007. Bladder cancer and exposure to water disinfection by-products through ingestion, bathing, showering, and swimming in pools. *Am J Epidemiol* 165(2): 148-156.

- Waage, A.S. et al. 1999. Detection of small numbers of *Campylobacter jejuni* and *Campylobacter coli* cells in environmental water, sewage, and food samples by a seminested PCR assay. *Appl Environ Microbiol* 65(4):1636-1643.
- Wade, T.J. et al. 2006. Rapidly measured indicators of recreational water quality are predictive of swimming-associated gastrointestinal illness. *Environ Health Perspect* 114(1):24-28.
- Wang, G., et al. 2007. Cancer risk assessment from trihalomethanes in drinking water. *Sci Tot Environ* 387:86-95.
- Watson, S.B. et al. 2003. Odours from pulp mill effluent treatment ponds: the origin of significant levels of geosmin and 2-methylisoborneol (MIB). *Chemosphere* 51:765-773.
- White, D.G. et al. 2000. Characterization of fluoroquinolone resistance among veterinary isolates of avian *Escherichia coli*. *Antimicrobial Agents and Chemotherapy* 44(10):2897-2899.
- White, D.G. et al. 2003. Antimicrobial susceptibilities of *Staphylococcus aureus* isolated from commercial broilers in northeastern Georgia. *Avian Diseases* 47:203-210.
- Whitman R.L. et al. 2003. Occurrence of *Escherichia coli* and Enterococci and *Cladophora* (Chlorophyta) in nearshore water and beach sand of Lake Michigan. *Appl Environ Microbiol* 69(8):4714-4719
- Wistrom, A.O. et al. 1996. A method for measuring haloform formation during wastewater chlorination. *Water Res* 30(12):3146-3151.
- World Health Organization (WHO). 1999. Toxic cyanobacteria in water: A guide to their public health consequences, monitoring and management. Edited by I. Chorus and J. Bartram. London, UK: E&FN Spon Publishers.
- World Health Organization (WHO). 2000. Monitoring bathing waters: A practical guide to the design and implementation of assessments and monitoring programmes (Chapter 9 – Approaches to microbiological monitoring; 10 – Cyanobacteria and algae). World Health Organization. Geneva, SWZ.
- World Health Organization (WHO). 2003. Guidelines for safe recreational water and environments: volume 1: coastal and fresh water. World Health Organization. Geneva, SWZ.
- World Health Organization (WHO). 2008a. Use of antimicrobials outside human medicine and resultant antimicrobial resistance in humans. <http://www.who.int/mediacentre/factsheets/fs268/en/index.html>.
- World Health Organization (WHO). 2008b. Drug-resistant Salmonella. <http://www.who.int/mediacentre/factsheets/fs139/en/index.html>.
- Xie, Y. et al. 2007. Determination of 2-methylisoborneol and geosmin produced by *Streptomyces* sp. and *Anabaena* PCC7120. *J Agric Food Chem* 55:6823-6828.
- Yoder, J.S. et al. 2004. Surveillance for waterborne-disease outbreaks associated with recreational water- United States, 2001-2002. *MMWR*, 2004: 53(No. SS-8)1-22.

Zhao, S. et al. 2005. Antimicrobial susceptibility and molecular characterization of avian pathogenic *Escherichia coli* isolates. *Vet Microbiol* 107:215-224.

ATTACHMENT B

Curriculum Vitae of Dr. Christopher M. Teaf

CURRICULUM VITAE

NAME: Christopher Morris Teaf

DATE OF BIRTH: 5 May 1953

PLACE OF BIRTH: Philadelphia, PA

HOME ADDRESS: 3527 Trillium Court
Tallahassee, FL 32312
(850) 668-4303

MARITAL STATUS: Married, two children

PROFESSIONAL POSITIONS: Associate Director
Center for Biomedical & Toxicological Research and Waste Management
Florida State University
2035 East Paul Dirac Drive
Suite 226 HMB
Tallahassee, FL 32310
(850) 644-5524 phone
(850) 574-6704 FAX
cteaf@fsu.edu email

President & Director of Toxicology
Hazardous Substance & Waste Management Research, Inc.
2976 Wellington Circle West
Tallahassee, FL 32309
(850) 681-6894 phone
(850) 906-9777 FAX
cteaf@hswmr.com email

EDUCATION: Ph.D. Toxicology, 1985
University of Arkansas for Medical Sciences (UAMS)
M.S. Biological Science, 1980
Florida State University (FSU)
B.S. Biology, 1975
Pennsylvania State University (PSU)

PROFESSIONAL MEMBERSHIPS: Florida Bar, Environmental and Land Use Law Section
National Association of Environmental Professionals
National Association of Underwater Instructors
Society of Toxicology
Society for Environmental Toxicology and Chemistry
Society for Risk Analysis

ACTIVITIES & HONORS:

Senior Editor for Human Risk Assessment, international journal *Human & Ecological Risk Assessment*, published by Taylor & Francis. 2004 to present (*also Editorial Board, 2000 to 2004*).

Editorial Board, international journal *Environmental Forensics*, Taylor & Francis Publ. 2006-present.

Advisory Board, Center for Terrorism & Public Health, FSU College of Medicine. 2001-present.

District 2 Local Emergency Planning Committee, Florida Emergency Response Commission, 1987 to present (*Vice Chair, 1991*).

Board of Directors, Dog Island Conservation District. 2002 to present (*Chair, 2004 to present*).

Science Advisory Board, 24th International Conference on Contaminated Soils, Sediments & Water, 2008. Amherst, MA.

Science Advisory Board, 23rd International Conference on Contaminated Soils, Sediments & Water, 2007. Amherst, MA.

Science Advisory Board, 22nd International Conference on Contaminated Soils, Sediments & Water, 2006. Amherst, MA.

Science Advisory Board, 21st International Conference on Contaminated Soils, Sediments & Water, 2005. Amherst, MA.

Editorial Board, international journal *Soil & Sediment Contamination*, CRC Press, LLC. 2002 to present.

Science Advisory Board, 20^h International Conference on Contaminated Soils, Sediments & Water, 2004. Amherst, MA.

Courtesy Professor, Department of Geology, Florida State University. 1999 to present.

Science Advisory Board, 19th International Conference on Contaminated Soils, Sediments & Water. Amherst, MA. 2003.

Science Advisory Board, 18th International Conference on Contaminated Soils, Sediments & Water. 2002. Amherst, MA.

Co-Director, NATO Advanced Research Workshop on Risk Assessment and Water Issues in Central Asia. 2000 to 2003. Almaty, Kazakhstan. 2000 to 2003.

Science Advisory Board, 17th International Conference on Contaminated Soils, Sediments & Water. 2001. Amherst, MA.

Advisory Board, Int'l Congress on Petrol. Contaminated Soils, Sediment & Water. London. 2000-2001.

Science Advisory Board, 16th International Conference on Contaminated Soils, Sediments & Water. 2000. Amherst, MA.

Assistant Scoutmaster, Troop 44. Tallahassee, FL. 1998 to 2004.

Editorial Board, *Environmental Toxicology & Chemistry*, published by CRC Press, LLC. 1997 to present.

Petroleum Underground Storage Tanks Technical Advisory Committee, Florida Department of Environmental Protection. 1996 - present.

Science Advisory Board, 15th International Conference on Contaminated Soils, Sediments & Water. Amherst, MA. 1998-1999.

Toxic Substances Advisory Council, Florida Department of Labor. 1990-2000 (*Chair, 1991-1996*).

Florida Bar, Environmental and Land Use Law Section Faculty and Steering Committee, 1996 - 1997.

Safety Committee, Florida Department of Environmental Protection, 1995 – 1997 (*advisor 1995-present*).

Florida Bar, Environmental and Land Use Law Section Faculty and Steering Committee, 1993 - 1994.

Technical Advisory Committee, MGP '95 - International Symposium on the Cleanup of Manufactured Gas Plant Sites. Prague, Czech Republic, 1994 to 1995.

Florida Comparison of Environmental Risks Project, 1994-1995 (*Human Health Co-chair*).

Landfill Technical Advisory Group, Florida Department of Environmental Protection, 1993-1994.

Benlate Health Effects Committee, Pesticide Review Council, Florida Department of Agriculture & Consumer Services, 1993-95.

Technical Advisory Committees for Budapest '92, Budapest '94, Warsaw '96, Warsaw '98, Prague 2000, Prague, 2003; 1st, 2nd, 3rd, 4th, 5th, and 6th International Symposia on Environmental Contamination in Central & Eastern Europe. 1991-2004.

Governor's Financial & Tech. Advisory Committee, Florida Dept. of Environ. Protection. 1986-1992.

Radon Measurement Specialist (*FL HRS Certification R1032*), 1989 to present.

Technical Referee: Human & Ecological Risk Assessment (1999-present); Nonlinearity in Biology, Toxicology and Medicine (2005-present); Ohio Journal of Science (1988-1992); Bulletin of Marine Sci. (1994-1998); Environmental Toxicology & Chemistry (1997-present); Environmental Biology of Fishes (1987-1990), Soil & Sediment Contamination (2002-present), Environmental Forensics (2006-present).

Director, Tallahassee Marathon/Half Marathon, 1989, 1990, 1991, 1995.

Outstanding Research Award, University of Arkansas Medical School, 1984.

Who's Who Among American College Students, 1983.

Graduation With Honors, Pennsylvania State University, 1975.

National Merit Scholar Program, 1971.

Eagle Scout, 1969 (Troop 1, Boy Scouts of America; Paoli, PA; Assistant Scoutmaster 1971 to 1975)

SUMMARY OF ACTIVITIES

Dr. Teaf received a BS (Biology) from Penn State and MS (Biological Science) from Florida State University. He earned a Ph.D. (Toxicology) from the University of Arkansas for Medical Sciences and conducted his research at the National Center for Toxicological Research. His experience includes positions of Associate Director, FSU Center for Biomedical & Toxicological Research and Hazardous Waste Management since 1983, and Director of Toxicology for HSWMR, Inc. since 1985 (President since 1989). From 1980 to 1982 he served as Research Staff for the Governor's Hazardous Waste Policy Advisory Council.

Areas of research and professional activity include: toxicological risk assessment for exposure to occupational and environmental chemicals under requirements of OSHA, CERCLA, SARA, RCRA, TSCA, and related state or federal legislation; modification of mutagenicity by environmental factors; effects of drugs and alcohol, male reproductive toxicology, aquatic toxicology, and waste management. Dr. Teaf has served as peer reviewer for: Human & Ecological Risk Assessment; Environmental Forensics; Nonlinearity in Biology, Toxicology and Medicine; Environmental Toxicology & Chemistry; Bulletin of Marine Science; Ohio Journal of Science; Chemosphere; Environmental Biology of Fishes, and reviews of research submitted to the Agency for Toxic Substances and Disease Registry (ATSDR). He is on Editorial Boards for Soil & Sediment Contamination and Environmental Forensics, and is Senior Human Risk Assessment Editor for Human & Ecological Risk Assessment. He has directed or conducted research for the U.S. Environmental Protection Agency (USEPA), U.S. Department of Agriculture, and several agencies in Florida, including: Department of Labor, Department of Environmental Protection (FDEP), Department of Health (FDOH), and Department of Community Affairs. He served as toxicologist to the Governor's Financial & Technical Advisory Committee (1986-1992), and for the state Landfill Technical Advisory Group (1993-1994). He presently serves as toxicologist for the state Petroleum Underground Storage Tanks Technical Advisory Committee. He was Co-chair of the Human Health Committee, Florida Comparison of Environmental Risks Project, a cooperative study funded by USEPA, FDEP and other agencies. He served as Chair of the Toxic Substances Advisory Council for the Florida Department of Labor, which implemented the Florida Right-to-Know Law. Dr. Teaf is toxicologist for and served as Vice-Chair of the District 2 Local Emergency Planning Committee (State Emergency Response Commission). From 1986-1989, he was liaison between the State University System of Florida Toxicological Research Coordinating Committee and FDEP. He presently is involved in activities in the U.S. and abroad related to preventing chemical terrorism. From 1998-2005, Dr. Teaf was actively involved in the statewide Contaminated Soils Forum and a technical subgroup, the Methodology Focus Group. He has served on graduate committees at Florida A&M University and Georgia Tech.

Chris has taught graduate and undergraduate courses at Florida State University, Florida A&M, the University of Florida, and Georgia Tech. He has organized and participated in seminars and training courses for many groups including the World Health Organization (WHO), USEPA, U.S. Department of Agriculture, ATSDR, FDOH, Florida Engineering Society, Florida Bar Association, FDEP, Florida Department of Education, Florida Chamber of Commerce, National Conference on Waste Exchange and Resource Reuse, National Hazardous Materials Training Center, and American Bar Association. He has served on Advisory Committees for MGP '95 (International Symposium on the Cleanup of Manufactured Gas Plants; in Prague), and for the six International Symposia on Environmental Contamination in Central & Eastern Europe (Budapest, Warsaw, Prague), among others.

Dr. Teaf has provided technical services on toxicology, environmental and human health risk assessment, waste management, recycling, water quality, and occupational health/safety, particularly regarding OSHA Hazard Communication Standard, Florida Right-to-Know Law and federal requirements under RCRA, CERCLA, and TSCA or related state regulations, including CERCLA cost allocations. He has provided services to the U.S. Attorney, and Florida State Attorney, as well as Attorneys General of Florida, Washington and Oklahoma. Dr. Teaf has testified on environmental issues, alcohol/drugs, occupational exposures, chemical health effects, and toxicological risk assessment to regulatory, legislative and judicial bodies.

SUMMARY OF PROFESSIONAL POSITIONS:

Associate Director
Center for Biomedical & Toxicological Research and Waste Management
Florida State University (Tallahassee, FL)
1983 to present

Director of Toxicology
Hazardous Substance & Waste Management Research, Inc. (HSWMR; Tallahassee, FL)
1985 to present (President, 1989 to present)

Advisory Board
Center for Terrorism and Public Health
Florida State University College of Medicine
2002 to present

Adjunct Faculty
College of Pharmacy & Pharmaceutical Sciences
Florida A & M University (Tallahassee, FL)
1998 to present

Board Member
Dog Island Conservation District
Franklin County, Florida
2002 to present (Chair, 2004 to present)

Board of Directors
Southern Waste Information Exchange (Tallahassee, FL)
2000 to present (Chair, 2000 to 2004)

Associate in Medicine
Program in Medical Sciences / FSU Medical School
Florida State University (Tallahassee, FL)
1995 to 2002

Toxicologist
Toxic Substances Advisory Council
Florida Department of Labor & Employment Security (Tallahassee, FL)
1990 - 1998 (Chairman, 1992 to 1998)

Toxicologist
Petroleum Underground Storage Tanks Advisory Committee
Florida Department of Environmental Protection (Tallahassee, FL)
1996 to present

Toxicologist
District 2 Local Emergency Planning Committee
Florida Department of Community Affairs (Tallahassee, FL)
1987 to present (Vice Chairman, 1991)

Toxicologist
Florida Landfill Technical Advisory Group
Florida Department of Environmental Protection (Tallahassee, FL)
1993 - 1994

Adjunct Faculty
Interdisciplinary Toxicology Program
University of Arkansas for Medical Sciences (Little Rock, AR)
1986 - 1989

Toxicologist
Governor's Financial and Technical Advisory Committee
Florida Department of Environmental Regulation (Tallahassee, FL)
1986 - 1992

Research Assistant
National Center for Toxicological Research and the University
of Arkansas for Medical Sciences (Little Rock, AR)
1983 - 1985

Associate Director
Southern Waste Information Exchange (Tallahassee, FL)
1981 - 1983

Research Staff
Florida Governor's Hazardous Waste Policy Advisory Council
1980 - 1982

Research Associate, Hazardous Waste Management Program
Florida State University (Tallahassee, FL)
1979 - 1983

Teaching/Research Assistant
Departments of Biological Science and Oceanography
Florida State University (Tallahassee, FL)
1976 - 1980

Animal Technician
Wyeth Laboratories (Philadelphia, PA)
1975 - 1976

TEACHING EXPERIENCE:

Air Quality (*Indoor/Outdoor*)
 Alcohol and Other Drugs
 Aquatic Toxicology
 Carcinogenesis and Carcinogens
 Comparative Vertebrate Morphology
 Dermal Toxicology
 Emergency Response and Contingency Planning
 Environmental Chemistry & Behavior of Chemicals
 General Zoology
 Hazardous Waste Management
 Health and Safety for Site Investigation and Response Personnel (*40 Hr and 8 Hr*)
 Ichthyology
 Inhalation Toxicology
 Industrial and Occupational Toxicology
 Metals Toxicology (*e.g., arsenic, lead, mercury, nickel, vanadium, cadmium, chromium*)
 Mutagenesis
 Nephrotoxicity
 Airborne Particulates
 Right-to-Know and Hazard Communication Laws (*OSHA and state*)
 Pesticide Toxicology
 Principles of Toxicology and Risk Assessment
 Radionuclides and Radiation Toxicology
 Reproductive Toxicology
 Research Diving Techniques
 Seafood Safety: Health Effects from Contaminated Products
 Toxicology of Solvents and Other Organic Hydrocarbons
 Toxicological Risk Assessment
 Toxicology and Management of Mixed Wastes (*Chemical/Radiological*)
 Toxicology for Physicians, Nurses, and Other Public Health Personnel

Courses listed represent those taught at Florida State University, University of Florida, University of North Florida, Florida A & M University, Georgia Institute of Technology and Tallahassee Community College. At Florida State University, Dr. Teaf has taught classes in the Departments of Biology, Chemistry, Engineering, Geology, Oceanography, and Urban & Regional Planning. Course coverage ranges from single day short courses to full semester graduate level and undergraduate level courses.

PUBLICATIONS AND SELECTED ABSTRACTS:

Weeks, N., D. Jones, C.M. Teaf, T. Lubozynski and Petrovich, M. 2007. RBCA and Conditional Closure as a Brownfields Tool. 10th Annual Florida Brownfields Conference & Exhibition. November, 2007. Orlando, FL.

Teaf, C.M., D.J. Covert and S. Kothur. 2007. Urban Polycyclic Aromatic Hydrocarbons (PAHs): A Florida Perspective. 23rd Annual International Conference on Soils, Sediments and Water. October, 2007. Amherst, MA.

Teaf, C.M. and B.L. Johnson. 2007. Ethics in Technical Publications: How Can the System Identify and Address Scientific Fraud. 23rd Annual International Conference on Soils, Sediments and Water. October, 2007. Amherst, MA. (*Also served as Session Moderator*).

Teaf, C.M., D.J. Covert, R. M. Coleman, M. Petrovich, R.S. Murali, and V. Yarina. 2007. Risk Issues and Background Evaluation for Arsenic in Soil at a Planned Residential Development. Contaminated Soils, Sediment & Water Volume 12: pp. 77-82. (*Also served as Session Moderator*).

Teaf, C.M. 2006. Disinfection byproducts: Benefits & limitations of existing drinking water guidelines. U.S. Environmental Protection Agency Federal-State Toxicology and Risk Analysis (FSTRAC) "Safe & Clean Water" Meeting. December, 2006. Clearwater, FL.

Khankhasayev, M., R. Herndon, J. Moerlins and C.M. Teaf. 2006. Overview of NATO CCMS Pilot Study on Environmental Decision-making for Sustainable Development in Central Asia. In: Vogtmann, H. and N. Dobretsov (eds.). Environmental Security and Sustainable Land Use – with special reference to central Asia. Springer Publishers, NATO Security through Science Series. 379 pp.

Coleman, R.M., C.M. Teaf, V.P. Cavey, D.J. Covert, S.N. Hughes, M. Marcus, M. McClure. 2006. Achievement of a Beneficial Use Designation for a Specialized High Volume Byproduct. Contaminated Soils, Sediments and Water. Volume 11, Chapter 3, pp. 27-48.

Teaf, C.M. 2006. Arsenic bioavailability workshop: Physical and chemical characteristics of soils which influence decisions regarding arsenic bioavailability and health risks. EPRI Arsenic Work Group. December, 2006. Tampa, FL

Teaf, C.M. and B.L. Johnson. 2006. Fraud and deception in publication of scientific research: Is there a solution? Human and Ecological Risk Assessment 12(4): 623-625.

Teaf, C.M. 2006. Forward to Volume 11. In: Kostecki, P.T., and E.J. Calabrese (eds.), Proceedings of the 21st Annual International Conference on Soils, Sediments and Water. Amherst, MA. Soil & Sediment Contamination, Volume 11, pp ix - x.

Teaf, C.M., B. Merkel, H.M. Mulisch, J.M. Kuperberg, and E. Wcislo. 2006. Industry, mining and military sites: Potential hazards and information needs. In: Schmoll, O. et al. (eds.), Protecting Groundwater for Health: Managing the Quality of Drinking-water Sources (Chapter 11). IWA Publishers, London, UK. pp. 309-336.

Teaf, C.M., B. Merkel, H.M. Mulisch, J.M. Kuperberg, and E. Wcislo. 2006. Industry, mining and military sites: Control and protection. In: Schmoll, O. et al. (eds.), Protecting Groundwater for Health: Managing the Quality of Drinking-water Sources (Chapter 23). IWA Publishers, London, UK. pp. 613-630.

White, M., D. Rountree, K. Geis, S. Kastury, K. Taylor, J. Curtis, D. Covert, and C.M. Teaf. 2006. Cascades Park: Remediation and Beneficial Recreational Redevelopment of a Former Manufactured

Gas Plant Site. 22nd Annual International Conference on Contaminated Soils, Sediments and Water. October, 2006. Amherst, MA.

C.M. Teaf, D.J. Covert, R. M. Coleman, M. Petrovich, R.S. Murali, and V. Yarina. 2006. Risk and Background Evaluation for Arsenic in Soil at a Planned Residential Development. 22nd Annual International Conference on Contaminated Soils, Sediments and Water. October, 2006. Amherst, MA.

Clark, B.S., P.T. Medico, F.J. Bermudez, M. Clewner, R.G. Wilkins, R.M. Coleman and C.M. Teaf. 2006. Beneficial Use of C&D Recovered Screen Material in Residential Applications: A Case Study. *Soil & Sediment Contamination* 11 (3): 355-360.

Teaf, C.M. 2006. A Toxicologist's Perspective on Industrial Exposures. *Workers First Watch*, Fall 2006. pp. 17-22. (Originally published as a paper in *Proceedings of the Annual Spring Conference & CLE for the Workers Injury Law & Advocacy Group*. May, 2006. Lake Buena Vista, FL.)

Teaf, C.M. and J.M. Kuperberg. 2006 Investigation and Information Collection for Contamination at Industrial Sites. In: Chorus, I. et al. (eds.), Monograph on Characterization and Mitigation of Surface Water Contamination. World Health Organization. In editorial review for publication in 2006.

Teaf, C.M. and J.M. Kuperberg. 2006 Critical Control Points and Prevention of Contamination for Industrial Sites. In: Chorus, I. et al. (eds.), Monograph on Characterization and Mitigation of Surface Water Contamination. World Health Organization. In editorial review for publication in 2006.

Teaf, C.M. and B.J. Tuovila. 2005. Indoor Air & Human Health: 21st Century Considerations. 8th Annual Florida Brownfields Conference. October, 2005. Jacksonville, FL.

Coleman, R.M., V.P. Cavey, D.J. Covert, S.N. Hughes, M. Marcus, M. McClure, and C.M. Teaf. 2005. Achievement of a Beneficial Use Designation for a Specialized High Volume Byproduct. 21st Annual International Conference on Contaminated Soils, Sediments and Water. October, 2005. Amherst, MA.

Clark, B.S., P.T. Medico, F.J. Bermudez, M. Clewner, R.G. Wilkins, R.M. Coleman and C.M. Teaf. 2005. Beneficial Use of C&D Recovered Screen Material in Residential Applications: A Case Study. 21st Annual International Conference on Contaminated Soils, Sediments and Water. October, 2005. Amherst, MA.

Teaf, C.M. Toxicology, Epidemiology & Public Health in the Courtroom. Florida Chamber of Commerce Summer School. July, 2005. Marco Island, FL.

Teaf, C.M. 2005. Science of Toxicology: Applications to Human Health Risk Assessment. Florida Association of Environmental Professionals Spring Conference. May, 2005. Jacksonville, FL.

Coleman, R.M., V.P. Cavey, D.J. Covert, S.N. Hughes, M. Marcus, M. McClure, and C.M. Teaf. 2005. Acquiring a Florida Beneficial Use Designation (BUD) for a High Volume Utility Byproduct. 42nd Annual Conference of the Air & Waste Management Association, Florida Section. November, 2005. St. Petersburg, FL.

Teaf, C.M., P.T. Medico, F.J. Bermudez, and B.S. Clark. 2005. Case Study: Beneficial Use of RSM in Residential Land Use Conditions. 15th Annual Materials Reuse and Recycling Conference. June, 2005. Key West, FL.

Teaf, C.M. 2005. Chemicals in the Workplace: How a Toxicologist Analyzes Exposure. 10th Anniversary Conference & CLE for the Workers Injury Law & Advocacy Group. April 10, 2005. Washington, DC.

Teaf, C.M. 2005. Critical Ecological Attributes and Historical Interstate Disputes Regarding the Apalachicola-Chattahoochee-Flint River System: A Panel Discussion. NATO Pilot Study Meeting: Tools and Approaches for Sustainability and Environmental Decision-Making in Central Asia. Wakulla Springs, FL. March, 2005.

Teaf, C.M. and B. W. Nocita. 2005. Site Investigation & Evaluation of Results: Assessment of Contamination and Risk. Technical Seminar for Hillsborough County Bar Association. February, 2005. Tampa, FL.

Teaf, C.M., H.V. Ritchie and R.M. Coleman. 2004. Status of the Science on Mold and Health Issues. Technical Seminar for First Coast Manufacturers Association. August, 2004. Jacksonville, FL.

Teaf, C.M., B.J. Tuovila, E.J. Zillioux, A. Shipp, G. Lawrence, C. Van Landingham. 2004. Nickel Carcinogenicity in Relation to the Health Risks from Residual Oil Fly Ash. *Human & Ecological Risk Assessment* 10(4): 665-682.

Teaf, C.M. and R.M. Coleman. 2004. Where Does the Science Stand on Mold Issues? Technical Seminar for White & Case Law Firm and for Marsh, Inc. January, 2004. Miami, FL.

Teaf C.M. and J.M. Kuperberg. 2004. Risk Assessment, Risk Management and Risk Communication: Principles and Applications. *In*: Teaf, C.M. et al. (eds.), *Risk Assessment as a Tool for Water Resources Decision-Making in Central Asia: Proceedings of a NATO Advanced Research Workshop*. Kluwer Academic Publishers, Dordrecht, Netherlands. NATO Science Series, Earth and Environmental Sciences 34: 1-15.

Teaf, C.M., B. Yessekin and M. Khankhasayev (eds). 2004. *Water Resources and Environmental Decision-Making in Central Asia: Proceedings of a NATO Advanced Research Workshop*. Kluwer Academic Publishers, Dordrecht, Netherlands. NATO Science Series, Earth and Environmental Sciences Volume 34. 324 pp.

Kuperberg, J.M., C.M. Teaf and H.V. Ritchie. 2003. Clean Up Approaches. *In*: Stapleton, R.M. (ed.), *Pollution A to Z*. Macmillan Reference Books, New York, NY.

Finley, B.L., T.J. Iannuzzi, N.D. Wilson, J.C. Kinnell, V. Craven, S. Lemeshow, C.M. Teaf, E.J. Calabrese and P.T. Kosteki,. 2003. The Passaic River Creel Angler Survey: Expert Panel Review, Findings and Recommendations. *Human and Ecological Risk Assessment* 9(3): 829-855.

DeMeo, R.A. and C.M. Teaf. 2003. Roles of the Attorney and Technical Consultants in Planning, Preparing and Presenting a Technical Case in Litigation. For National Registry of Environmental Professionals (NREP) and the International Society of Technical and Environmental Professionals (INSTEP). Orlando, FL. November, 2003.

Teaf, C.M. 2003. Overview of Risks Posed by Biological and Chemical Contaminants in Drinking Water. NATO Pilot Study Meeting: Tools and Approaches for Sustainability and Environmental Decision-Making in Central Asia. Almaty, Kazakhstan. November, 2003.

Aertgeerts, R., J. Colford and C.M. Teaf. 2003. Potential Health Effects of Saline Waters: Toxicology & Epidemiology. *Proceedings of the Sixth International Symposium on Environmental Contamination in Central and Eastern Europe, Prague, Czech Republic*. September, 2003. (*Also served as Session Moderator*).

Teaf, C.M., D.J. Covert and R.M. Coleman. 2003. Risk Assessment Applications Beyond Baseline Risks and Clean-up Goals. *Soils & Sediment Contamination* 12(4): 497-506. July, 2003.

Teaf, C.M. 2003. Chemical Agents Relevant to Terrorism Against Food Supplies and Water. 2003. Statewide Health Department Video Conference on Chemical Terrorism. For Florida State University College of Medicine and Florida Department of Health. June, 2003.

Teaf, C.M., D.J. Covert and R.M. Coleman. 2003. Arsenic in Florida: An Update on the SCTL Process. For Air & Waste Management Association. Pensacola, FL. May, 2003.

Teaf, C.M. 2003. Toxicology and adverse health impacts from the consumption of highly saline water supplies. World Health Organization Public Health Initiatives Meeting. Copenhagen, Denmark. April, 2003.

Teaf, C.M., D.J. Covert and R.M. Coleman. 2002. Risk Assessment Applications in Atypical Circumstances. 18th Annual International Conference on Contaminated Soils, Sediments and Water. Amherst, MA. October, 2002.

Teaf, C.M. and J.M. Kuperberg. 2002. Development of Health-based Water Criteria for Environmental Chemicals: Theory and Application. NATO Advanced Research Workshop: Risk Assessment as a Tool for Environmental Decision-Making in Central Asia. Almaty, Kazakhstan. September, 2002.

Teaf, C.M. 2002. Risk Assessment & Risk Management: Goals and Applications. NATO Advanced Research Workshop on Water Resources and Environmental Decision-Making in Central Asia. Almaty, Kazakhstan. September, 2002.

Teaf, C.M., J.M. Kuperberg, H.V. Ritchie and B.J. Tuovila. 2002. Vulnerability of Water Systems to Chemical Attack. In: R. Aertgeerts (ed.), Prevention and Management of Substance Terrorism Against Water Services. World Health Organization Monograph. 85 pp.

I. Petrisor, M. Stefanescu, S. Dobrota, A. Voicu and C.M. Teaf. 2002. Microbiota Involved in Vegetation of Sulphidic Mine Tailings: Greenhouse Experiments. 18th Annual International Conference on Contaminated Soils, Sediments and Water. Amherst, MA. October, 2002.

Teaf, C.M., J.M. Kuperberg, and H.V. Ritchie. 2002.. Toxicology and the Environmental Behavior of Selected Chemical Terrorism Agents. Environmental Response Team Training, Florida Department of Environmental Protection, Division of Law Enforcement. Camp Blanding, FL. April, 2002.

Teaf, C.M., H.V. Ritchie, and J.M. Kuperberg. 2002. Chemical Terrorism: Properties and Toxicological Effects of Selected Agents. Conference on Terrorism in the New Millenium. Tampa, FL. April, 2002.

Petrisor, I., K. Komnitsas, I. Lazar, J.M. Kuperberg and C.M. Teaf. 2002. Studies on Biosorption of Heavy Metals from Mine Effluents. Proceedings of the 12th Annual West Coast Conference on Contaminated Soil, Sediment and Water. San Diego, CA. March, 2002.

Teaf, C.M., J.M. Kuperberg, I.G. Petrisor, E. Wcislo, R. Kucharski, T. Hazen and A. Tien. 2001. Bioremediation of a Petroleum-contaminated Site in Southern Poland: Phase 1 – Site Characterization and Risk Assessment. Proceedings of the First International Congress on Petroleum Contaminated Soils, Sediments & Water. London, UK, August 2001. (*Also served as Session Moderator*)

Petrisor, I.G., S. Dobrota, A. Voicu, M. Stefanescu, C. Teaf, M. Kuperberg. 2001. Emulsifying Activity of Several Good Hydrocarbon Degrading Bacterial Strains. Proceedings of the First International Congress on Petroleum Contaminated Soils, Sediments & Water. London, UK, August 2001.

Kuperberg, J.M., C.M. Teaf, I.G. Petrisor, A. Worsztynowicz, M. Adamski, M. Heitkamp, T. Hazen and A. Tien. 2001. Bioremediation of a Petroleum-contaminated Site in Southern Poland: Phase 2 – Biopile Construction and Operation. Proceedings of the First International Congress on Petroleum Contaminated Soils, Sediments & Water. London, UK, August 2001.

Teaf, C.M. 2001. Risk Assessment Principles and Practice in the US. Planning Seminar on Introduction of the Rapid Environmental and Health Risk Assessment Protocol for the Lower Danube River. World Health Organization and Italian Ministry of the Environment. Sofia, Bulgaria, January, 2001.

Petrisor, I.G., S. Dobrota, I. Lazar, A. Voicu, M. Stefanescu, M. Kuperberg, and C. Teaf,. 2001. Field Deployment of Oily Sludge Bioremediation in Romania. Proceedings of the 2001 International Containment & Remediation Technology Conference and Exhibition. Orlando, FL.

Teaf, C.M. 2000. Toxic Effects of Chlorinated and Nonchlorinated Solvents. *In*: Williams, P. et al. (eds.), Principles and Practices of Toxicology: Industrial and Environmental Applications. John Wiley & Sons, Inc., New York, NY.

Donkin, S.R., D.L. Ohlson and C.M. Teaf. 2000. Properties and Effects of Metals. *In*: Williams, P. et al. (eds.), Principles and Practices of Toxicology: Industrial and Environmental Applications. John Wiley & Sons, Inc., New York, NY.

Teaf, C.M. and P. Middendorf. 2000. Mutagenesis and Genetic Toxicology. *In*: Williams, P. et al. (eds.), Principles and Practices of Toxicology: Industrial and Environmental Applications. John Wiley & Sons, Inc., New York, NY.

Smoak, J.M., M. Brenner, M.S. Allen, C.M. Teaf and B.J. Tuovila. 2000. Elevated radium in Florida lakes. Proceedings of the American Geophysical Union meeting, San Francisco, CA. December, 2000.

Abdel-Rahman, M.S., H. Brown, B.G. Callahan, R. Di Giulio, C. Metcalfe, C. Teaf, and R. Yang. 2000. Peer Review Panel Report for the Fox River Human and Ecological Risk Assessments. Association for Environmental health & Sciences, Amherst, MA. June, 2000.

Leeper, D.A., M.H. Kelly, M. Brenner, J.M. Smoak, W. Kenney, T.J. Whitmore, M.S. Allen, C.L. Schelske, B.J. Tuovila and C.M. Teaf. 2000. Radium-226 in Florida lakes: bioaccumulation and environmental risks. North American Lake Management Society 2000 Annual Meeting. Miami, FL. November, 2000.

Khan, A., S. Pack and C.M. Teaf. 2000. Dilution Attenuation Factors for RBCA at Miami International Airport. Proceedings of the Air & Water Resource Association Annual Conference. Miami, FL. November, 2000.

Kuperberg, J.M., C.M. Teaf, S. Dushenkov, C. Negri, R. Kucharski and A. Sas-Nowosielska. 2000. Phytoremediation – risk-based evaluation. Proceedings of the Fifth International Symposium on Environmental Contamination in Central and Eastern Europe, Prague, Czech Republic. September, 2000. (*Also served as Session Moderator*).

Rygiel, P.R. and C. M. Teaf. 2000. Disparate Information Obtained From Separate Analytical Techniques for Arsenic Determination in Groundwater: Implications for Risk Assessment. Proceedings of the Fifth International Symposium on Environmental Contamination in Central and Eastern Europe, Prague, Czech Republic. September, 2000. (*Also served as Session Moderator*).

Teaf, C.M., B.J. Tuovila, J.M. Kuperberg, D.A. Leeper, M.H. Kelly, M. Brenner, J.M. Smoak, M.S. Allen. 2000. Potential Health and Environmental Risks Associated with ²²⁶Radium in Florida Lakes. Proceedings of the Fifth International Symposium on Environmental Contamination in Central and Eastern Europe, Prague, Czech Republic. September, 2000. (*Also served as Session Moderator*).

Sellers, L., S. Emery, J. Wojtecki, and C.M. Teaf. 2000. Managing Sites with Chemical Issues: Evaluation and Presentation of Risk Information. Florida Chamber of Commerce 14th Annual Environmental Permitting Summer School. Marco Island, FL, July, 2000.

Kuperberg, J.M., K. Henderson, K.F.A. Soliman, C.M. Teaf and M.G. Kolta. 1999. Time of administration affects organophosphate pesticide-induced alterations in body temperature. *Toxicological Sciences* 48 (1-S): 187.

Teaf, C.M. 1999. Risk Assessment for Chemical and Radioactive Materials: Some Characteristics, Similarities and Differences. In: Choppin, G.R. and M.K. Khankhasayev (eds.), Chemical Separation Technologies and Related Methods of Nuclear Waste Management – Applications, Problems and Research Needs. Proceedings of the NATO Advanced Study Institute, held in Dubna, Russia, May, 1998. Kluwer Academic Publishers, Dordrecht, Netherlands. pp. 261-272.

Teaf, C.M., R.G. Cooper and J.J. McNally. 1999. Arsenic associated with golf course operations: on-site and off-site risk considerations for soils and water. Annual Meeting of the Society for Risk Analysis. Atlanta, GA. December, 1999.

Teaf, C.M. 1999. Can regulatory agency actions accommodate hormesis, adaptation and mechanisms of toxicity? *Biological Effects of Low Level Exposures* 8 (1): 19-21.

Teaf, C.M. 1999. Risk-Based Corrective Action (RBCA) at Miami International Airport. 1999 Hazardous Waste Management Conference. Miami, Florida. May, 1999.

Teaf, C.M. 1999. Toxicology and Risk Assessment: Using the Expert Witness. 1999 Annual Meeting, National Association of Legal Assistants. Tampa, FL. July, 1999.

Fleming, L.E., J.A. Bean, I.K. Stabile and C.M. Teaf. 1999. Evaluation of epidemiologic studies on the human health effects associated with hazardous waste incineration facilities. In: Roberts, S.M., J. Bean and C.M. Teaf (eds.), Hazardous Waste Incineration: Evaluating the Human Health and Environmental Risks (Chapter 10). CRC Press, Boca Raton, FL. pp. 253-282.

Teaf, C.M., I.K. Stabile and P. Moffat. 1999. Characterization of potential emissions from hazardous waste incinerators and related facilities. In: Roberts, S.M., J. Bean and C.M. Teaf (eds.), Hazardous Waste Incineration: Evaluating the Human Health and Environmental Risks (Chapter 2). CRC Press, Boca Raton, FL. pp. 23-50.

Teaf, C.M., I.K. Stabile, P. Moffat and J.M. Kuperberg. 1999. Physical, chemical and toxicological properties of potential emissions from hazardous waste incinerators and related facilities. In: Roberts, S.M., J. Bean and C.M. Teaf (eds.), Hazardous Waste Incineration: Evaluating the Human Health and Environmental Risks (Chapter 3). CRC Press, Boca Raton, FL. pp. 51-85.

Teaf, C.M. 1999. Toxicology and Risk Assessment Considerations Concerning Chlorinated Solvents in Groundwater. For Florida Department of Health and Agency for Toxic Substances and Disease Registry (ATSDR). Wakulla Springs, FL. May, 1999.

Teaf, C.M. 1998. Session Moderator's Summary for Human Health Risk Assessment. *Human and Ecological Risk Assessment* 4(3): 671-674.

Tonner-Navarro, L.E., J.L. Phelps, S.M. Roberts and C.M. Teaf. 1998. Methods for Developing Risk-based Cleanup Goals for Complex Mixtures. *Human and Ecological Risk Assessment* 4(3): 721-736.

Goodheart, G.F., C.M. Teaf and M.J. Manning. 1998. Coupling Risk-Based Remediation with Innovative Technology. *Pollution Engineering* 30(5): 48-50.

Kerger, B.D. and C.M. Teaf. 1998. Validating airborne chemical exposure assessment for juries: two case studies involving chlorinated hydrocarbons in groundwater. Society for Risk Analysis, Risk Science and Law Specialty Group. Phoenix, AZ. December, 1998.

Wcislo, E., J.M. Kuperberg and C.M. Teaf. 1998. Risk Assessment as a Tool for Guiding Phytoremediation Activities: A Case Study for the Bytom Site in Poland. Proceedings of the 1998 Fourth International Symposium on Environmental Contamination in Central and Eastern Europe, Warsaw, Poland. September, 1998. *(Also served as Session Moderator)*.

Teaf, C.M., D.J. Rillstone and M.J. Manning. 1998. Evaluation of Risks Posed by the Use of Contaminated Groundwater for the Manufacture of Concrete. Proceedings of the 1998 Fourth International Symposium on Environmental Contamination in Central and Eastern Europe, Warsaw, Poland. September, 1998. *(Also served as Session Moderator)*.

Teaf, C.M. 1998. Toxicology & Health-Based Risk Assessment: Applications in the Workplace. Proceedings of the National Aeronautics & Space Administration (NASA) Occupational Health Conference. Orlando, FL. August 24, 1998.

Teaf, C.M. 1998. Escambia Wood Treatment Facility: Investigations and Public Health Issues. For Florida Department of Health and Agency for Toxic Substances and Disease Registry (ATSDR). Annual Meeting of the Florida Environmental Health Association. Cocoa Beach, FL. June, 1998.

Teaf, C.M. 1998. Risk Assessment for Chemicals & Radionuclides: Some Characteristics, Similarities and Differences. NATO Advanced Study Institute of Chemical Separation Technologies. Dubna, Russia. May, 1998.

Teaf, C.M. 1998. Toxicology & Health-Based Risk Assessment: Applications in the Workplace. American Industrial Hygiene Association. March, 1998.

Teaf, C.M. 1998. Implementation of Risk-Based Corrective Action (RBCA) Under Different State and Federal Cleanup Programs. Florida Bar Association, Miami, FL. January, 1998.

Kuperberg, J.M., M.G. Kolta, R. Kuharski, E. Wcislo, L. Gray and C.M. Teaf. 1997. Use of Risk-Based Approaches to Guide Environmental Remediation Research in Poland. Society of Environmental Toxicology and Chemistry, San Francisco, CA. November, 1997.

Teaf, C.M., R.M. Coleman, M.J. Manning and D.J. Covert. 1997. Site-specific Modification of the Standard ASTM-Tiered Approach for a Complex Site in Florida. Society of Environmental Toxicology and Chemistry, San Francisco, CA. November, 1997.

Teaf, C.M. 1997. Brownfield Development, Petroleum Sites and Risk-Based Corrective Action (RBCA): Where Have We Been and Where Are We Going? Annual Meeting of the Environmental and Land Use Law Section of the Florida Bar. Naples, FL. August, 1997.

Teaf, C.M. 1997. How Clean is Clean Enough? The Use of Risk Assessment Techniques in Brownfields Projects. Air & Waste Management Association. Gatlinburg, TN. August, 1997.

Teaf, C.M., S.M. Roberts and E. Akin (*co-panelists*). 1997. RBCA Cleanup Strategies for Brownfields Properties. Presented at Conference entitled The Challenges of Eastward Ho!: RBCA, Brownfields and Environmental Justice. Miami, FL. June, 1997.

Teaf, C.M., M.J. Manning and G.F. Goodheart. 1997. Use of Tiered Approaches to Cleanup Objectives at Illinois Brownfields Sites. Presented at Annual Meeting of the National Association of Environmental Professionals. May, 1997. *(Also served as Session Moderator)*.

Kuperberg, J.M., M.G. Kolta, E. Wcislo and C.M. Teaf. 1997. The application of risk-based approaches to the evaluation and remediation of a refinery site in Poland. Annual Meeting of the Society of Toxicology. Cincinnati, OH. March, 1997.

Teaf, C.M. 1997. Risk-Based Corrective Actions at RCRA Sites. University of Florida TREEO Hazardous Waste Management Course. Ft. Lauderdale, FL. January 16, 1997.

Teaf, C.M., R.M. Coleman, M.J. Manning, D.J. Covert and J.L. Phelps. 1996. Multipathway Human Health Risk Assessment Concerning Air Emissions from Combustion of Orimulsion Fuel. Society for Environmental Toxicology and Chemistry. Washington, D.C. November 21, 1996.

Teaf, C.M. 1996. Risk-based corrective action: principles and practices. Florida Association of Environmental Professionals. Ft. Lauderdale, FL. October 18, 1996.

Teaf, C.M. 1996. Risk assessment practices and guidelines in the southern states. 1996 Environmental Expo. Tampa, FL. October 3, 1996.

Teaf, C. M and I.K. Stabile. 1996. Site-specific evaluation of potential public health risks: A case study of a Florida Superfund site. Annual Meeting of the Florida Public Health Association. Sarasota, FL. September 18, 1996.

Kuperberg, J.M., M. Kolta and C.M. Teaf. 1996. Phytoremediation: Potential for application in Central and Eastern Europe (C&EE). Proceedings of the 1996 Third International Symposium on Environmental Contamination in Central and Eastern Europe, Warsaw, Poland. September, 1996. (*Also served as Session Moderator*).

Kuperberg, J.M., E. Wcislo and C.M. Teaf. 1996. Application of risk-based approaches to the remediation of a Polish refinery site. Proceedings of the 1996 Third International Symposium on Environmental Contamination in Central and Eastern Europe, Warsaw, Poland. September, 1996. (*Also served as Session Moderator*).

Lillich, R.B., C.M. Teaf, D.H. Stinson, W. Green, and R.D. La Belle. 1996. Environmental Dilemmas Facing Eastern Europe and the Former Soviet Union. *Journal of Transnational Law & Policy* 5(2):1-42.

Teaf, C.M. 1996. Toxicology and CERCLA Risk Assessments: Common Approaches & Applications. American Bar Association, Section on Litigation. Vail, CO. February 19, 1996.

Roberts, S.M. and C.M. Teaf. 1995. Filling in the gaps: Addressing toxicity potential not typically included in risk assessments. Presented at 1995 Conference on State of Practice of Risk Assessment in Human Health and Environmental Decision Making. Tallahassee, FL. December 13-14, 1995. (*Also served as Session Moderator*).

Kuperberg, J.M., C.M. Teaf and M.G. Kolta. 1995. Relative Toxicity as a Function of Environmental Availability. *Land Contamination and Reclamation* 3(4):3-4 to 3-5. (*Also served as Session Moderator*).

Teaf, C.M., R.C. Herndon and D.M. Haney. 1995. Risk Assessment: Principles and Applications to Historical Manufactured Gas Plant Sites. *Land Contamination and Reclamation* 3(4):3-21 to 3-23. (*Also served as Session Moderator*).

Wilson, I., I. Stabile and C.M. Teaf. 1995. Behavior modification by means of targeted education in high-risk raw oyster consumers. Submitted to *Florida Journal of Public Health*.

Teaf, C.M. 1995. Principles and Applications of Health-based Risk Assessment. *Keep Florida Beautiful*, Fall/Winter Edition 1995: pp. 26-29.

Teaf C.M. 1995. Human Health and Environmental Risks Associated with Contaminated Military Sites. In: R.C. Herndon et al. (eds.), *Cleanup of Former Soviet Military Installations: Identification and*

Selection of Environmental Technologies for Use in Central and Eastern Europe. pp. 31-44. Springer-Verlag, New York, NY.

Stabile, I.K. and C.M. Teaf. 1995. International Conference on Health Effects of Hazardous Wastes, Atlanta, GA. June, 1995. (*Also served as Session Moderator*).

Teaf, C.M. 1995. Evaluating Potential Risks to the Public from a Superfund Site: Case Study. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). Annual Meeting of the Florida Environmental Health Association. Cocoa Beach, FL. May 11, 1995.

Teaf, C.M. 1995. Medical Toxicology and Public Health Issues related to Florida Hazardous Waste Sites. For FSU Program in Medical Sciences, Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). February, 1995.

Teaf, C.M. 1995. Worker Health and Safety Requirements. RCRA Inspector Training Institute, for U.S. Environmental Protection Agency and Florida Department of Environmental Protection. Orlando, FL. January 24-26, 1995.

Teaf, C.M. 1995. Weighing the Odds: An Overview and Perspective on Environmental Risk Assessments. 11th Annual Advanced Environmental Permitting Short Course, Tallahassee, FL. January 20, 1995.

Teaf, C.M. and E.B. Jones. 1995. Risk Assessments for Contaminated Sites in Relation to Real Estate Transactions. Quarterly Meeting of the Florida Envir. Assessors Assoc., Tallahassee, FL. January, 1995.

Harbison, R.D., C.J. Borgert and C.M. Teaf. 1995. Placental Metabolism of Xenobiotics. *In*: Sastry, B.V.R., Placental Toxicology. pp. 213-238. CRC Press, Boca Raton, FL.

Teaf, C.M., I.K. Stabile and J. Winter. 1994-1995. Seafood Poisoning: Biological and Nonbiological Effects of Contaminated Fish and Shellfish from the Gulf of Mexico. For U.S. Environmental Protection Agency Gulf of Mexico Program and Florida Department of Health and Rehabilitative Services. Tallahassee, FL, Corpus Christi, TX, Lafayette, LA, Biloxi, MS, Daytona Beach, FL and Huntsville, AL. (March, May, July, August, October, November, 1994 and June, 1995).

Teaf, C.M. 1994. Risk Assessment Approaches, Procedures and Goals. Florida Bar Environmental and Land Use Law Section Workshop, Tampa, FL. October, 1994. (*Also served as Workshop Co-Chair*).

Biczó, I., Sz. Böröcz, P. Richter, C.M. Teaf and M. Kuperberg. 1994. Developing a Chemical Safety Management System for Chlorine. Second International Symposium on Environmental Contamination in Central and Eastern Europe, Budapest, Hungary. September, 1994. (*Also served as Session Moderator*).

Teaf, C.M. and J.M. Kuperberg. 1994. Toxicology and Public Health Considerations in the Development of Cleanup Criteria for Contaminated Sites. Proceedings of the 1994 Second International Symposium on Environmental Contamination in Central and Eastern Europe, Budapest, Hungary. September, 1994. (*Also served as Session Moderator*).

Teaf, C.M. 1994. Health-based Risk Assessment and the Remediation of Contaminated Sites: Principles and Applications. For North Atlantic Treaty Organization (NATO) and the Joint Center for Hungarian-American Environmental Research, Studies & Exchanges (CHAERSE). Visegrad, Hungary. June, 1994.

Teaf, C.M. and D. Grant. 1994. Using Risk Assessment as a Remedial Option. Annual Florida Environmental Resources Conference. Orlando, FL. May 26, 1994.

Teaf, C.M. 1994. Hazardous Materials: Toxicology & Emergency Response. Medical Seminars on Public Health Implications of Hazardous Materials Incidents. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). Tallahassee, FL. March, 1994.

Teaf, C.M. 1994. Toxicology and Public Health: Agents, Exposures & Effects. Medical Seminars on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). Sarasota, Miami and Tallahassee, FL. January, February and March, 1994.

Winter, J.M. and C.M. Teaf. 1993. Benlate: a medical and scientific controversy. J. Florida Medical Association 80(6):400-402.

Teaf, C.M. and I.K. Stabile. 1993. Hazardous Materials: Toxicology and Emergency Response. Medical Seminars on Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). Pensacola and Tallahassee, FL. November, 1993.

Teaf, C.M. and I.K. Stabile. 1993. Toxicology and public health issues: Florida hazardous waste sites. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry (ATSDR). Pensacola, Ft. Lauderdale, Tampa and Sarasota, FL. March, April, May, October and November, 1993.

Teaf, C.M. 1993. Manufactured Gas Plant Sites: Application of Risk Assessment in the Development of Remedial Target Concentrations for Manufactured Gas Plant Sites. American Gas Association 1993 Distribution & Transmission Conference, Orlando, FL. May 19, 1993.

Winter, J.M. and C.M. Teaf. 1993. Physician Education in Diagnosis and Treatment of Diseases Related to Environmental and Occupational Exposures to Hazardous Waste. ATSDR International Conference on Health Effects of Hazardous Wastes, sponsored by ATSDR, Atlanta, GA. May, 1993.

Herndon, R.C., P.I. Richter, J.E. Moerlins, J.M. Kuperberg, and C.M. Teaf. 1993. The 1992 International Symposium on Environmental Contamination in Central and Eastern Europe: A Forum for Technology Transfer. International Conference on Health Effects of Hazardous Wastes, sponsored by ATSDR, Atlanta, GA. May, 1993. *(Also served as Session Moderator)*.

Teaf, C.M. 1992. Toxicological Risk Assessment in the Application of Health-based Target Concentrations to Site Remediation. Proceedings of the 1992 International Symposium on Environmental Contamination in Central and Eastern Europe, Budapest, Hungary. October, 1992. *(Also served as Session Moderator)*.

Haney, D.M., R.C. Herndon and C.M. Teaf. 1992. Historical Manufactured Gas Plant Sites: Use of Risk Assessment in the Development of Remedial Target Concentrations. Engineering Technical Note ENV-92-4-1, pp. 14. Operating Section, American Gas Association. November, 1992.

Teaf, C.M. and I.K. Stabile. 1992. Toxicology and public health issues: Florida hazardous waste sites. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami and Orlando, FL. April, May and October, 1992.

Teaf, C.M. and I.K. Stabile. 1992. Site exercise: Evaluating potential risks from a hazardous waste site. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami and Orlando, FL. April, May and October, 1992.

I.K. Stabile and C.M. Teaf. 1992. Differential diagnosis and treatment of acute and chronic exposure to chemicals at hazardous waste sites. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami and Orlando, FL. April, May and October, 1992.

Stabile, I.K. and C.M. Teaf. 1992. A case study in environmental medicine: trichloroethylene exposure. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami and Orlando, FL. April, May and October, 1992.

Stabile, I.K. and C.M. Teaf. 1992. A case study in environmental medicine: lead exposure. Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami and Orlando, FL. April, May and October, 1992.

Teaf, C.M. 1991. Risk Assessment and the Development of Remedial Targets for Contaminated Sites. Presented at Annual Environmental Engineering Conference, University of Florida. November, 1991.

Teaf, C.M. 1991. The Importance of Risk Assessment to Industry. Presented at Ninth Annual Carolina Conference on Environmental Law and Technology. Greenville, SC. June 1991.

Teaf, C.M. 1991. Dioxin Toxicity Revisited: Critical Characteristics and Current Applications. Presented at 1991 Mississippi Environment Meeting. Jackson, MS. May 1991.

Teaf, C.M. 1991. Toxicology and the Derivation of Maximum Contaminant Levels (MCLs). Presented at Annual Meeting of the Underground Injection Practices Council. Mobile, Alabama. February, 1991.

Teaf, C.M. and I.K. Stabile. 1990-1991. Toxicology, public health and patient response to health advisories. Physician Training Course on Public Health Implications of Environmental Toxicants. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa, Orlando, Ft. Lauderdale and Tallahassee, FL. March, July, November, 1990; April, September, 1991.

I.K. Stabile and C.M. Teaf. 1990-1991. Pediatric aspects of environmental health. Physician Training Course on Public Health Implications of Environmental Toxicants. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa, Orlando, Ft. Lauderdale and Tallahassee, FL. March, July, November, 1990; April, September, 1991.

I.K. Stabile and C.M. Teaf. 1990-1991. Clinical evaluation of suspected environmental and occupational illness. Physician Training Course on Public Health Implications of Environmental Toxicants. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa, Orlando, Ft. Lauderdale and Tallahassee, FL. March, 1990; July, 1990; November, 1990; April, 1991; and September, 1991.

Teaf, C.M. and I.K. Stabile. 1990-1991. A case study for methylene chloride exposure: therapy and patient education. Physician Training Course on Public Health Implications of Environmental Toxicants. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa, FL. March, 1990.

Teaf, C.M. and I.K. Stabile. 1990-1991. A case study for organophosphate pesticide exposure: therapy and patient education. Physician Training Course on Public Health Implications of Environmental Toxicants. Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa, Orlando, Ft. Lauderdale and Tallahassee, FL. March, 1990; July, 1990; November, 1990; April, 1991; and September, 1991.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1990. Depression of glutathione in male reproductive tissues and potentiation of EMS-induced germ cell mutagenesis by 1,2-dibromoethane. *Teratogenesis, Carcinogenesis, Mutagenesis* 10(6):428-438.

Stabile, I.K., C.M. Teaf and R.D. Harbison. 1989. Non-neuronal cholinergic and carnitineric systems in mammalian tissues. *Contemporary Reviews in Obstetrics and Gynecology* 1:253-260.

Teaf, C.M. 1989. Development of Site Rehabilitation Levels (SRLs) at contaminated sites. State Program Managers Annual Meeting, Association of State and Territorial Solid Waste Management Officials. Tucson, AZ. November, 1989.

Teaf, C.M., I.K. Stabile and R.D. Harbison. 1989. Potentiation of in vivo male germ cell mutagenesis by agents which deplete glutathione. European Society of Human Reproduction and Embryology. Copenhagen, Denmark. June, 1989.

Herndon, R.C., J.E. Moerlins, C.M. Teaf, V.W. Lambou and J.D. Koutsandreas. 1988. Hazardous waste from small quantity generators in the United States. Hazardous Waste: Detection, Control, Treatment. M Elsevire Science Publishers, Amsterdam, Netherlands.

Bishop, J.B., C.M. Teaf and V.E. Campbell. 1988. Assessment of fetal death among in utero progeny of Fischer 344 rats after gavage treatment of males with 2-AAF or 4-AAF. *In*: Ashby, J. et al. (eds.). Evaluation of Short-term Tests for Carcinogens: Report of the International Programme on Chemical Safety's Collaborative Study of in Vivo Assays, Volume 2. pp. 215-225. Cambridge University Press, Cambridge, UK.

Bishop, J.B., C.M. Teaf and M.F. Salamone. 1988. Assessment for abnormality in head shape of epididymal sperm from Fischer 344 male rats treated by gavage with 2-AAF or 4-AAF. *In*: Ashby, J. et al. (eds.). Evaluation of Short-term Tests for Carcinogens: Report of the International Programme on Chemical Safety's Collaborative Study of in Vivo Assays, Volume 2. pp. 258-260. Cambridge University Press, Cambridge, UK.

Teaf, C.M., R.C. Herndon and R.M. Coleman. 1988. Development of toxicant profiles for gasoline and other selected environmental contaminants. Proc. 1st Annual Environmental Toxicology Research Conference. Tallahassee, FL. June, 1988.

Teaf, C.M., R.C. Herndon and R.M. Coleman. 1988. Modeling relative risks from multiple routes of exposure to toxicants. Proc. 1st Annual Environment Toxicology Research Conference. Tallahassee, FL. June, 1988.

Teaf, C.M. 1988. Development and applicability of site rehabilitation levels. Florida Pollutant Storage Tanks Program. Annual Meeting. Orlando, FL. June, 1988.

Teaf, C.M. 1988. Methyl bromide toxicology and the assessment of risk for low-level exposures. U.S. Department of Agriculture Animal and Plant Health Inspection Service Annual Meeting. Lumberton, NC. April, 1988.

Teaf, C.M. 1988. Toxicology and human health effects of methyl bromide. Methyl Bromide Workshop. Florida Department of Agriculture and Consumer Services. La Belle, FL. March, 1988.

Jones, E.B., C.M. Teaf, J.E. Moerlins and R.C. Herndon. 1987. Current developments in the role of waste exchanges for satisfying waste minimization requirements. Annual Meeting of the Florida Society of Environmental Analysts. Orlando, FL. October, 1987.

Freeman, R.W. and C.M. Teaf. 1987. General principles of toxicology and fire safety precautions. Arson Conference. Florida Office of the State Fire Marshal. Orlando, FL. September, 1987.

Herndon, R.C., C.M. Teaf and J.E. Moerlins. 1987. Hazardous waste from United States small quantity generators. World Conference on Hazardous Waste. Budapest, Hungary. October, 1987.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1987. Depression of glutathione in male reproductive tissues and potentiation of EMS-induced germ cell mutagenesis by L-buthionine sulfoximine. *Teratogenesis, Carcinogenesis, Mutagenesis* 7(6):497-514.

Teaf, C.M. and T.C. Lewis. 1987. Seasonal occurrence of multiple caudal spines in the Atlantic stingray, Dasyatis sabina. *Copeia* 1987(1):224-227.

Bishop, J.B., C.M. Teaf and B. Booshan. 1987. Assessment of fetal death rate among in utero progeny of B6C3 and CD-1 mice after subcutaneous injection of males with butyl benzylphthalate. *Environmental Mutagenesis* 9(Suppl.8):15.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1986. Glutathione and the potentiation of ethyl methanesulfonate (EMS)-induced germ cell mutagenesis by ethylene dibromide (EDB) and buthionine sulfoximine (BSO). *Environmental Mutagenesis* 8(6):85.

Gandy, J., C.M. Teaf, F.A. Adatsi, R.C. James and R.D. Harbison. 1986. Dependence of male reproductive toxicity on germ cell stage. In: Fujii, T. and P.M. Adams (eds.). Functional Teratogenesis: Proceedings of the Fourth International Congress of Toxicology. Teikyo Univ. Press, Tokyo, Japan.

Teaf, C.M. 1986. Toxicological aspects of risk assessment. In: Proceedings of the 1985 National Governors' Association Conference on Toxics in Groundwater. Committee on Energy and Environment, National Governors' Association. Washington, DC.

Moerlins, J.E., R.C. Herndon and C.M. Teaf. 1985. Hazardous Waste Management Guide for Florida Hospitals. Florida Hospital Association, pp. 180.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1985. Germ cell mutagenesis and glutathione (GSH) depression in reproductive tissue of the F-344 rat induced by ethyl methanesulfonate. *Mutation Research* 144:93-98.

Teaf, C.M. 1985. Mutagenesis. In: P.L. Williams and J. Burson (eds.). Industrial Toxicology: Safety and Health Applications in the Workplace. Van Nostrand-Reinhold Co., pp. 502.

James, R.C. and C.M. Teaf. 1985. Carcinogenesis. In: P.L. Williams and J. Burson (eds.). Industrial Toxicology: Safety and Health Applications in the Workplace. Van Nostrand-Reinhold Co., pp. 502.

Teaf, C.M., J.B. Bishop and R.D. Harbison, 1985. Ethylene dibromide (EDB) and buthionine sulfoximine (BSO) potentiation of ethyl methanesulfonate (EMS)-induced dominant lethal mutations. *Toxicologist* 5(1):140.

Teaf, C.M., R.W. Freeman and R.D. Harbison. 1984. Cocaine-induced hepatotoxicity: Lipid peroxidation as a possible mechanism. *Drug and Chemical Toxicology* 7(4):383-396.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1984. Alteration of ethyl methanesulfonate (EMS) dominant lethal effects in Fischer 344 rats with buthionine sulfoximine. *Genetics* 107(3):105.

Domon, O.E., C.M. Teaf, B. Booshan and J.B. Bishop. 1984. Assessment of the fetal death rate among *in utero* progeny of B6C3 and CD-1 mice after subcutaneous injection of males with

(diethylhexyl)phthalate (DEHP). International Phthalic Acid Ester Conference, Surrey, UK. September, 1984.

Teaf, C.M., J.B. Bishop and R.D. Harbison. 1984. Effect of cadmium and buthionine sulfoximine (BSO) on glutathione levels in the male reproductive tract. *Toxicologist* 4(1):80.

Teaf, C.M., R.M. Freeman and R.D. Harbison. 1983. Cocaine-induced hepatotoxicity: Lipid peroxidation as a possible mechanism. *Toxicologist* 3(1):95.

Herndon, R.C., J.E. Moerlins and C.M. Teaf. 1983. Waste exchange and resource reuse. *Resource Recycling* 2(4):22-26.

Teaf, C.M., D.C. White and R.C. Herndon. 1981. The nature of hazardous waste: chemistry, health effects and management strategies. Annual Meeting, Florida Bar Environmental Law Section.

Teaf, C.M. 1981. Occurrence and distribution of selected vertebrates, vegetation and endangered species in Florida. In: Fernald, E.A. (ed.). Atlas of Florida, FSU Foundation, Inc.

Teaf, C.M. and T.C. Lewis, 1981. Seasonal occurrence of multiple caudal spines in the Atlantic Stingray, Dasyatis sabina. *ASB Bulletin* 28(2):55.

Teaf, C.M. 1981. Current status of state and federal regulations governing management of hazardous wastes in Florida. Florida Division Annual Conference, American Chemical Society. May, 1981.

Teaf, C.M. 1980. Phylum Chordata: The Vertebrates (a laboratory guide to vertebrate dissection). In: Tschinkel, W.R. and P.C. Hayward, An Introduction to Animal Diversity, Burgess Publishing Company, Inc.

Teaf, C.M. 1980. Tidally-oriented movement patterns of the Atlantic stingray, Dasyatis sabina. *ASB Bulletin* 27(2): 6.

TECHNICAL REPORTS AND CONTRACTS:

Use of Glass from Cathode Ray Tubes in Concrete Products. For Southern Waste Information eXchange and Florida Department of Environmental Protection. 2007-2008. (ongoing).

Development and Performance of RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Protection. 2005-present. (ongoing; Contract HW-527).

Development and Performance of RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 2000-2005. (Contract HW-432).

The Florida Hazardous Waste Regulation Program: History and Positive Impacts on Florida's Environment. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 1998 (Contract HW-330).

RCRA Compliance Assistance Training Course for the Fiberglass Manufacturing Industry. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 1996-1999 (Contract HW-330).

Development and Performance of RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 1996-1999 (Contract HW-330).

Leon County Emergency Management Plan for Hazardous Materials at Fixed Facilities. For Leon County Board of Commissioners, Leon County Division of Emergency Management and Florida Department of Community Affairs. 1996-1997.

RCRA Compliance Assistance Training Course for the Automotive Maintenance & Repair and Auto Paint & Body Industries. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 1993-1996 (Contract HW-242).

Development and Performance of RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Protection (Bureau of Waste Planning and Regulation). 1993-1996 (Contract HW-242).

Toxicological Support Services and Public Health Training for Physicians and Environmental Health Personnel. For Florida Department of Health & Rehabilitative Services. 1995-1997.

Public Education and Human Health Aspects of Seafood Safety. For U.S. Environmental Protection Agency Gulf of Mexico Program. 1995-1996.

Potential Human Health Impacts and Environmental Effects from Hazardous Waste Incinerators and Related Facilities. For Florida Department of Environmental Protection, and Florida Center for Solid & Hazardous Waste Management. 1994-1995 .

Seafood Safety: Toxicology and Microbial Issues Related to Contaminated Products. For U.S. Environmental Protection Agency Gulf of Mexico Program. 1994-1995.

Development of Site Rankings for Petroleum Contamination Sites. For Florida Department of Environmental Regulation. 1992-1994 (Contract HW-204).

Toxicological Support Services: Reviews of Site-Specific Risk Assessments and the Development of Acceptable Exposure Guidelines for Hazardous Waste Sites (e.g., CERCLA, State-lead). For Florida Department of Environmental Regulation (Division of Waste Mgmt). 1991-1993 (Contract WM-445).

Development and Performance of Specific RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Regulation (Bureau of Waste Planning and Regulation). 1992-1993 (Contract HW-218).

Leon County Emergency Management Plan for Hazardous Materials at Fixed Facilities. For Leon County Board of Commissioners, Leon County Division of Emergency Management and Florida Department of Community Affairs. 1992-1993.

Technical Evaluation and Regulatory Review of RCRA Facility Closure Risk Assessments. For Florida Department of Environmental Regulation (Bureau of Waste Planning and Regulation). 1991-1992 (Contract HW-195).

Development and Performance of Specific RCRA-related Personnel Health and Safety Training Activities. For Florida Department of Environmental Regulation (Bureau of Waste Planning and Regulation). 1991-1992 (Contract HW-196).

Quality Assurance Review and Evaluation for Petroleum Contamination Sites Database. For Florida Department of Environmental Regulation (Division of Waste Mgmt). 1990-1991 (Contract GC-128).

Leon County Emergency Management Plan for Hazardous Materials at Fixed Facilities. For Leon County Board of Commissioners, Leon County Division of Emergency Management and Florida Department of Community Affairs. Final Report. May, 1991.

Toxicological Support Services: Reviews of Site-Specific Risk Assessments and the Development of Acceptable Exposure Guidelines for Hazardous Waste Sites (e.g., CERCLA, State-lead). For Florida Department of Environmental Regulation (Division of Waste Mgmt). 1989-1991 (Contract WM-331).

Technical Evaluation and Regulatory Review of RCRA Facility Closure Risk Assessments. For Florida Department of Environmental Regulation (Bureau of Waste Planning and Regulation). 1989-1991 (Contract HW-141).

Development of Site Rankings for Petroleum Contamination Sites. For Florida Department of Environmental Regulation. 1989-1992 (Contract HW-126).

Development of Site Rehabilitation Levels per Chapter 17-70, F.A.C. For Florida Department of Environmental Regulation. 1989-1990 (Contract HW-135).

Risk Assessment for Establishment of Acceptable Contaminant Levels in Environmental Media. (Toxicant Profiles for 8 compounds). For Florida Department of Environmental Regulation. 1988-1989 (Contract WM-318).

Florida Non-Hazardous Industrial-Waste-Disposal Facilities. Two volume final report prepared for the U.S. EPA on the characteristics and proximity of such facilities to sensitive environments.

Monitoring Guidance for Municipal-Waste-Combustion-Ash Disposal. #EPA/600/X-90/267A&B. 1990.

Development of Automated Systems for Assessing Subtitle D Facility Locations. #EPA/600/X-88/423. December, 1988.

Wet Environments RCRA Subtitle D Monitoring Guidance Parts I-IV. #EPA/600/X-88/403A,B,C,&D. December, 1988.

Characterization of Municipal Waste Combustion Ash Disposal Practices and Facilities (Preliminary Summary of Seventy-Two Municipal Waste Combustion Ash Disposal Facilities). #EPA/600/X-88/360. October, 1988.

Facility/Site Description of Pinellas County Solid Waste Management Complex Pinellas County, Florida. For U.S. Environmental Protection Agency. March, 1988.

Facility/Site Description of Southeast Hillsborough County Sanitary Landfill Hillsborough County, Florida. For U.S. Environmental Protection Agency. March, 1988.

Leon County Comprehensive Emergency Management Plan for Hazardous Materials. For Leon County Board of Commissioners, Division of Emergency Management and Florida Department of Community Affairs. Final Report. March, 1989.

1989 Annual Report of the State University System Toxicological Research Coordinating Committee. For the Florida Legislature and Office of the Governor. Final Report. March, 1989.

Risk Assessment for Establishment of Acceptable Contaminant Levels in Environmental Media. (Toxicant Profiles for 15 compounds) For Florida Department of Environmental Regulation. 1988-1989 (Contract WM-251).

Toxicological Support Services: Reviews of Site-Specific Risk Assessments and the Development of Acceptable Exposure Guidelines for Hazardous Waste Sites under CERCLA. For Florida Department of Environmental Regulation (Division of Waste Cleanup). 1988-1989 (Contract WM-223).

Technical Evaluation and Regulatory Review of RCRA Facility Closure Risk Assessments. For Florida Department of Environmental Regulation (Bureau of Environmental Permitting). 1988-1989 (Contract HW-98).

Locational Survey of Underground Storage Tank Facilities in Florida for Use in the State's Intergraph Data Base System. For Florida Department of Environmental Regulation (Bureau of Waste Management, Storage Tank Regulation Section). Final Report. January, 1989.

Development of Monitoring Guidance with an Emphasis on Wet Environments. For U.S. Environmental Protection Agency. 1987-1994 (ongoing).

Toxicological Support Services: Reviews of Site-Specific Risk Assessments and the Development of Acceptable Exposure Guidelines for Hazardous Waste Sites under CERCLA. For Florida Department of Environmental Regulation (Division of Waste Management). 1987-1988 (Contract WM-223).

Environmental Audit and Development of a Waste Management Plan. For Baptist Medical Center. Jacksonville, Florida. 1987.

1987 Annual Report of the State University System Toxicological Research Coordinating Committee. For the Florida Legislature and Office of the Governor. Final Report. March, 1987.

Relative Risks from Multiple Routes of Exposure to Toxicants Found in Water Supplies: A Review and Evaluation. For Florida Department of Environmental Regulation (Bureau of Groundwater Protection). Final Report. November, 1987.

Toxicological Support Services: Reviews of Site-Specific Risk Assessments and the Development of Acceptable Exposure Guidelines for Hazardous Waste Sites (e.g., CERCLA, State-lead). For Florida Department of Environmental Regulation (Bureau of Operations). 1986-1987 (Contract WM-186).

Risk Assessment for Establishment of Acceptable Contaminant Levels in Environmental Media. (Toxicant Profiles for 15 compounds) For Florida Department of Environmental Regulation. 1985-1987. Final Reports. November, 1987.

Toxicological Data Analysis and Risk Assessment For Facilities Identified During the Stationary Compression Vessels Notification Program. For Florida Department of Environmental Regulation. Final Report. December, 1986.

1986 Annual Report of the State University System Toxicological Research Coordinating Committee. For the Florida Legislature and Office of the Governor. Final Report. March, 1986.

Manual for Training Courses in Toxicology: Hazardous Waste Field Activities. For Florida Department of Environmental Regulation. Final Report. June, 1986.

Hazardous Waste Management Guide for Automotive Repair and Supply Shops. For Florida Automotive Services Council. Final Report. May, 1986.

Manual for Training Course on Emergency Response and Contingency Planning. For the U.S. Environmental Protection Agency. 1986.

Risk Assessment for Determination of Acceptable Levels of Selected Toxic Compounds in Florida Groundwater. (Toxicant Profiles for 28 compounds). For Florida Department of Environmental Regulation, Florida Board of Regents and Florida Department of Health and Rehabilitative Services. 1983-1987. Final Revised Reports. November, 1987.

Training Manual - Seminar on the Florida Right-to-Know Law and Management of Hazardous Wastes by Hospitals. For Florida Hospital Association. 1985.

Hazardous Waste Management Guide for Florida Hospitals. For Florida Hospital Association. 1985.

Proceedings of the Second National Conference on Waste Exchange and Resource Reuse. 1985.

1985 Annual Report of the State University System Toxicological Research Coordinating Committee. For the Florida Legislature and Office of the Governor. Final Report, March, 1985.

Alternative Technologies for Hazardous Waste Management in Florida. For the Florida Institute of Government and the Florida Legislature. 1984.

Toxicological Evaluation Manual: Risk Assessment and Safety Determination for Chemical Exposures. For the Florida Department of Health and Rehabilitative Services. 1984.

Development of a National Training Center for Hazardous Materials Management: A Concept Paper. For State of Florida and U.S. Environmental Protection Agency. 1983.

Leon County Biological Relationships. For Leon County Board of Commissioners and City of Tallahassee. 1983.

Hazardous Waste Issues in Florida: A Legislative Monograph. For Florida House of Representatives and Florida Senate. 1983.

Proceedings of the First National Conference on Waste Exchange and Resource Reuse. 1983.

Southern Waste Information Exchange Catalog. Published quarterly 1982 to present.

East Pass Lagoon Water Quality Study: Phases I, II, III. For Okaloosa Co. Bd. of Commissioners. 1982.

Siting of a Hazardous Waste Management Facility: An Overview. 1982.

Hazardous Waste: A Management Perspective. Final Report of the Governor's Hazardous Waste Policy Advisory Council to the Florida Legislature. 1981.

Hazardous Materials Contingency Plans for Local Governments. For U.S. Environmental Protection Agency. 1981.

Feasibility Study for Establishing a Florida Waste Exchange. For Florida Department of Environmental Regulation. 1981.

An Analysis of the Local, State, and Federal Authority and Responsibility for Hazardous Materials Incidents in Leon County, Florida. For Leon County Board of Commissioners. 1980.

Hazardous Waste Incidents in the State of Florida (Volume II). For Florida Department of Environmental Regulation. 1980.

Assessment of the Economic Impact of Florida's Proposed Hazardous Waste Control Program on Selected Industries. For Florida Department of Environmental Regulation. 1979.

Hazardous Waste Incidents in the State of Florida (Volume I). For Florida Department of Environmental Regulation. 1978.

Hazardous Waste Survey for the State of Florida. For Florida Department of Environmental Regulation. 1977.

SELECTED CONFERENCES AND TRAINING COURSES:

Personnel Health and Safety Refresher Training Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, FL. April, 2008.

Weeks, N., Jones, D., Teaf, C.M., Lubozynski, and Petrovich, M. 2007. RBCA and Conditional Closure as a Brownfields Tool. 10th Annual Florida Brownfields Conference & Exhibition. November, 2007. Orlando, FL.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, FL. April, 2007.

Teaf, C.M. and L. Gray. 2007. Principles, Terminology & Applications of Toxicology in a University Setting. For Environmental Health & Safety Department, Florida State University, Tallahassee, FL. April, 2007

Teaf, C.M. 2006. Disinfection byproducts: Benefits & limitations of existing drinking water guidelines. U.S. Environmental Protection Agency Federal-State Toxicology and Risk Analysis (FSTRAC) "Safe & Clean Water" Meeting. December, 2006. Clearwater, FL.

Teaf, C.M. 2006. Arsenic bioavailability workshop: Physical and chemical characteristics of soils which influence decisions regarding arsenic bioavailability and health risks. EPRI Arsenic Work Group. December, 2006. Tampa, FL

Teaf, C.M. and P.H. Burress. 2006. Principles, Terminology & Applications of Toxicology in a University Setting. For Environmental Health & Safety Department, Florida State University, Tallahassee, FL. July, 2006.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, FL. April, 2006.

Teaf, C.M., P.T. Medico, F.J. Bermudez, and B.S. Clark. 2005. Case Study: Beneficial Use of RSM in Residential Applications:. 15th Annual Materials Reuse and Recycling Conference. June, 2005. Key West, FL.

Teaf, C.M. 2005. Chemicals in the Workplace: How a Toxicologist Analyzes Exposure. 10th Anniversary Conference & CLE for the Workers Injury Law & Advocacy Group. April 2005. Washington, DC.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, FL. April, 2005.

Khankhasayev, M., R. Herndon, J. Moerlins and C. Teaf. 2005. Overview of NATO CCMS Pilot Study on Environmental Decision-making for Sustainable Development in Central Asia. Presented at NATO Advanced Research Workshop (ARW) on Environmental Security and Sustainable Land Use of the Mountain and Steppe Territories of Mongolia and Altai: Present Day Role of Traditions and Science. Presented at NATO ARW, Barnaul, Russia, October 2004.

Teaf, C.M., H.V. Ritchie and R.M. Coleman. 2004. Status of the Science on Mold and Health Issues. Technical Seminar for First Coast Manufacturers Association. August, 2004. Jacksonville, FL.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. November, 2004.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. March, 2004.

Teaf, C.M. and R.M. Coleman. 2004. Where Does the Science Stand on Mold Issues? Technical Seminar for White & Case Law Firm and for Marsh, Inc. January, 2004. Miami, FL.

NATO Pilot Study Meeting on Human Health Issues and Tools for Disease Prevention in Central Asia. Almaty, Kazakhstan. November, 2003. (*Served as Session Moderator*).

Sixth International Symposium on Environmental Contamination in Central and Eastern Europe. Prague, Czech Republic. September, 2003. (*Session Moderator and Steering Committee Member*).

OSHA Refresher Course for University Health and Safety Activities (8 hours). For Florida State University. Tallahassee, Florida. June 11, 2003.

Statewide Video Conference on Chemical Terrorism. For Florida State University College of Medicine and Florida Department of Health. June 10, 2003.

Personnel Health and Safety Refresher Course (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. May 7, 2003.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. January 8, 2003.

NATO Advanced Research Workshop: Risk Assessment as a Tool for Environmental Decision-Making in Central Asia. Almaty, Kazakhstan. September, 2002. (*Served as Co-Director and Editor of Proceedings*).

Safety Concerns: Awareness for the Field Inspector. For Florida Department of Environmental Protection. St. Petersburg, Florida. August, 2002

Vulnerability of Municipal and Private Water Systems to Chemical Attack. Presented at World Health Organization Restricted Expert Consultation on Prevention and Management of Substance Terrorism Against Water Services. Copenhagen, Denmark. November, 2001.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. May, 2001.

Safety Concerns: Awareness for the Field Inspector. For Florida Department of Environmental Protection. St. Petersburg, Florida. August, 2001

Planning Seminar on Introduction of the Rapid Environmental and Health Risk Assessment for the Lower Danube River. For the World Health Organization and Italian Ministry of the Environment. Sofia, Bulgaria. January, 2001.

Evaluating and Communicating Health Risks. For Holland & Knight Annual Environmental Seminar Series. Orlando, Florida. October, 2000.

Fifth International Symposium on Environmental Contamination in Central and Eastern Europe. Prague, Czech Republic. September, 2000. (*Session Moderator and Steering Committee Member*).

14th Annual Environmental Permitting Summer School. Managing sites with chemical issues: Evaluation and presentation of risk information. For Florida Chamber of Commerce. Marco Island, Florida. July, 2000.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Ft. Myers, Florida. August, 1999.

Toxicology and Risk Assessment: Using the Expert Witness. 1999 Annual Meeting, National Association of Legal Assistants. Tampa, FL. July, 1999.

Risk-Based Corrective Action at Miami International Airport. 1999 Hazardous Waste Management Conference. Miami, Florida. May, 1999.

Toxicology and Risk Considerations for Chlorinated Solvents. For Florida Department of Health & Rehabilitative Services. Wakulla Springs, Florida. May, 1999.

Fourth International Symposium on Environmental Contamination in Central and Eastern Europe. Warsaw, Poland. September, 1998. (*Session Moderator and Steering Committee Member*).

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Ft. Myers, Florida. July, 1998.

Risk Assessment Principles & Practice: Application to the Polish Environment. For Institute for Ecology of Industrial Areas and U.S. Department of Energy. Katowice, Poland. January, 1997.

Control of Reportable Diseases in Florida: A Public-Private Partnership. For Florida Department of Health & Rehabilitative Services. Tallahassee, Florida. October 24, 1996.

Third International Symposium on Environmental Contamination in Central and Eastern Europe. Warsaw, Poland. September, 1996. (*Session Moderator and Steering Committee Member*).

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. June 6, 1996.

Environmental Toxicology for Physicians *and* Medical History-taking in the Evaluation of Environmental or Occupational Disease. For Florida Department of Health & Rehabilitative Services. Sebring, Florida and Vero Beach, Florida. May 10 and May 15, 1996.

Control of Reportable Diseases in Florida: A Public-Private Partnership. For Florida Department of Health & Rehabilitative Services. Tallahassee, Florida. April 17, 1996.

RCRA Compliance Technical Assistance Training Courses. For Florida Department of Environmental Protection. Tallahassee, Florida (Eleven locations in Florida during period February to August, 1996).

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. April 2, 1996.

American Bar Association Section of Litigation, 8th Annual Environmental Litigation Midyear Meeting. Vail, Colorado. February 15-16, 1996.

12th Annual Environmental Permitting Short Course. For Florida Chamber of Commerce. Orlando, Florida. January 17-18, 1996.

1995 Conference on State of Practice of Risk Assessment in Human Health and Environmental Decision Making. Tallahassee, FL. December 13-14, 1995. (*Session Moderator and Steering Committee Member*).

Risk Assessments, Audits & Other Compliance Management Tools. Environmental Compliance & Risk Management Seminar, Florida Chamber of Commerce. Tampa, FL. November, 1995.

Air Contamination in Central and Eastern Europe: What We Have Seen Since the Iron Curtain was Pulled Back. For Carolinas Air Pollution Control Association. Myrtle Beach, SC. October, 1995.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. September 27, 1995.

MGP'95 International Symposium and Trade Fair on the Cleanup of Manufactured Gas Plants. Prague, Czech Republic. September, 1995. (*Session Moderator and Steering Committee Member*).

Seafood Poisoning: Biological and Nonbiological Effects of Contaminated Fish and Shellfish from the Gulf of Mexico. For U.S. Environmental Protection Agency and Florida Department of Health and Rehabilitative Services. Tallahassee, Florida; Corpus Christi, Texas; Lafayette, Louisiana; Biloxi, Mississippi; Daytona Beach, Florida and Orange Beach, Alabama. March 26, May 22, July 15, August 3, August 18, October 13, October 18, November 11, 1994 and June 6, 1995.

Required Personnel Health and Safety Refresher Course for Hazardous Waste Activities (8 hours). For Levine•Fricke. Tallahassee, Florida. August 14, 1995.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. May 18, 1995.

Worker Health and Safety Requirements. For U.S. Environmental Protection Agency RCRA Inspector Training Institute. Orlando, Florida. January 24-26, 1995.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Ft. Myers, Florida. January 12, 1995.

RCRA Personnel Health and Safety Training Course (40 hours). For Florida Department of Environmental Protection. Orlando, Florida. December 5-9, 1994.

Training Course in Principles of Toxicology, Risk Assessment and Risk Communication. Conducted for Florida Department of Health and Rehabilitative Services. Jacksonville, Florida. November, 1994.

Risk Assessments: Understanding Their Strengths and Weaknesses (*Session Co-Chair*). For the Florida Bar Association, Environmental and Land Use Law Committee. Tampa, Florida. October 10, 1994.

Training Course for RCRA Inspectors. For Florida Department of Environmental Protection. Tallahassee, Florida. October 4-6, 1994.

Second International Symposium on Environmental Contamination in Central and Eastern Europe. Budapest, Hungary. September 20-23, 1994. (*Session Moderator and Steering Committee Member*).

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Protection. Tallahassee, Florida. August 23, 1994.

Regulations, Environmental Considerations and Safety Aspects of Petroleum Sites under Chapter 17-770. For Florida Department of Environmental Protection. Tallahassee, Florida. June 7-8, 1994.

Toxicology & Risk Assessment Training Course. Conducted for Florida Department of Environmental Protection. Tallahassee, Florida. May 18-19, 1994.

Seminars for Physicians and Medical Personnel on Public Health Implications of Toxic Materials and Hazardous Waste Sites. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Miami, Sarasota and Tallahassee, Florida. March 3, March 11, March 12, March 25 and March 30, 1994.

Required Personnel Health and Safety Refresher Course for Hazardous Waste Activities (8 hours). For Levine•Fricke. Tallahassee, Florida. March 7, 1994.

Training Course in Principles of Toxicology, Risk Assessment and Risk Communication. Conducted for Florida Department of Health & Rehabilitative Services. Jacksonville, Florida. February 3-4, 1994.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. January 8, 1994.

RCRA Personnel Health and Safety Training Course (40 hours). For Florida Department of Environmental Regulation. Orlando, Florida. December 6-10, 1993.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. July 8, 1993.

Personnel Health and Safety Refresher Course for RCRA Activities (8 hours). For Florida Department of Environmental Regulation. Orlando, Florida. May 20, 1993.

Medical Seminars on Public Health Implications of Hazardous Waste Sites. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Pensacola, Ft. Lauderdale and Miami, Florida. March, April, May, November, 1993.

Training Course in Principles of Toxicology, Risk Assessment and Risk Communication. Conducted for Florida Department of Health and Rehabilitative Services. Ft. Myers, Florida. January 28-29, 1993.

Personnel Health and Safety Refresher Course for Hazardous Waste Activities (8 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. December 9, 1992.

RCRA Personnel Health and Safety Training Course (40 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. November 16-20, 1992.

Required Personnel Health and Safety Refresher Course for Hazardous Waste Activities (8 hours). For Levine•Fricke. Tallahassee, Florida. November 18, 1992.

First International Symposium on Environmental Contamination in Central and Eastern Europe. Budapest, Hungary. October 21-24, 1992. (*Session Moderator and Steering Committee Member*).

Training Course in Principles of Toxicology and Risk Assessment. For Florida Department of Health and Rehabilitative Services. Orlando, Florida. July 23-24, 1992.

RCRA Personnel Health and Safety Required Refresher Course (8 hours). For Florida Department of Environmental Regulation. Orlando and Tallahassee, Florida. May 29, June 23 and September 15, 1992.

Training Course in Principles of Toxicology and Risk Assessment. Conducted for Florida Department of Health and Rehabilitative Services. Tallahassee, Florida. June 18-19, 1992.

General Principles of Toxicology and Risk Assessment (3 Graduate Semester Hours). For Department of Biological Science, Florida State University. Tallahassee, Florida. January to April, 1992.

Medical Seminar on Community Public Health Implications of Hazardous Waste Sites. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Rockledge, Miami, and Orlando, Florida. April 10, April 24, May 28 and October 22, 1992.

Personnel Health & Safety Refresher Course for Hazardous Waste Activities (8 hours). For Florida Department of Environmental Regulation. Orlando, Tallahassee, FL. April, October, November, 1991.

RCRA Personnel Health and Safety Training Course (40 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. January 4-8 and September 16-20, 1991.

Physician Training Course on Public Health Implications of Environmental Toxicants. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. St. Petersburg and Orlando, Florida. April 12, September 13 and September 25, 1991.

Supervisors Site Health and Safety Training Course for Hazardous Waste Activities (8 hours). For Levine•Fricke. Tallahassee, Florida. June 18, 1991.

Required Personnel Health and Safety Refresher Course for Hazardous Waste Activities (8 hours). For Levine•Fricke. Tallahassee, Florida. June 17, 1991.

Training Course on General Principles of Toxicology and Risk Assessment. Conducted for Florida Department of Health and Rehabilitative Services. Orlando, Florida. May 1-2 and May 21-22, 1991.

Training Course on Mixed Radioactive and Hazardous Waste. For Florida Department of Environmental Regulation. Tallahassee, Florida. October, 1990.

Physician Training Course on Public Health Implications of Environmental Toxicants. For Florida Department of Health and Rehabilitative Services and Agency for Toxic Substances and Disease Registry. Tampa and Tallahassee, Florida. March 15 and July 24, 1990.

Mercury Contamination in Florida: Impacts and Solutions. Conducted by FSU Center for Biomedical & Toxicological Research, sponsored by 19 federal and state agencies. June 20-21, 1990.

Environmental Toxicology & Epidemiology: A Practical Approach for Local Health Officials and Physicians. For National Association of County Health Officials (NACHO), Agency for Toxic Substances and Disease Registry, and Florida Department of Health. Ft. Lauderdale, FL. July, 1990.

RCRA Personnel Health and Safety Training Course (40 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. June 4-8, 1990.

RCRA Personnel Health and Safety Required Refresher Course (8 hours). For Florida Department of Environmental Regulation. Tallahassee, Florida. May 31, 1990.

RCRA Personnel Health and Safety: Basic Training Course (40 hours). For Florida Department of Environmental Regulation. Orlando, Florida and Tallahassee, FL. April 24-28 and December 4-8, 1989.

Groundwater Investigations and Application to Assessment of Risks. For Florida Department of Environmental Regulation. Tallahassee, Florida. June 6-7, 1989.

Risk Assessment & Decision-Making Training. Conducted for Florida Department of Environmental Regulation and U.S. Environmental Protection Agency. Tallahassee, Florida. August 30-31, 1988.

Training Course on General Principles of Toxicology and Risk Assessment. Conducted for Florida Department of Health and Rehabilitative Services. Tallahassee, Florida. September 2-3, 1987 and Orlando, Florida. October 27-28, 1987.

Hazardous Materials Training (Risk Assessment; Hazardous Materials Contingency Plans). Conducted for National Hazardous Materials Training Center, Little Rock, Arkansas. October 28-30, 1986.

Training Course on Toxicology: Hazardous Waste Field Activities. Conducted for the Florida Department of Environmental Regulation and the Florida Department of Health and Rehabilitative Services. Tallahassee, Florida. June 25-26, 1986 and Orlando, Florida. October 8-9, 1986.

Training Course on Emergency Response and Contingency Planning. Conducted for the U.S. Environmental Protection Agency. Atlanta, Georgia. March, 1985; March, 1986.

Training Course for the Holmes Regional Medical Center: Employee Toxic Substances Right-to-Know Program. Conducted for the Holmes Regional Medical Center. Melbourne, Florida. 1986.

Training Seminar on the Florida Right-to-Know Law and Management of Hazardous Wastes by Hospitals. Conducted for the Florida Hospital Association (FHA). Orlando, Florida. August, 1986.

Training Seminar on the Florida Right-to-Know Law and Management of Hazardous Wastes by Hospitals. Conducted for the Florida Hospital Association (FHA). Orlando, Florida. November, 1985.

Hazardous Waste Disposal and Hazardous Substances Right-to-Know Regulations. Conducted for Florida Hospital Engineers Association. Orlando, Florida. April 4, 1986.

Second National Conference on Waste Exchange & Resource Reuse. Tallahassee, Florida. March, 1985.

Training Course on the OSHA Hazard Communication Standard. Orlando, Florida. February, 1985.

Symposium: Alternative Technologies for Waste Management. Tallahassee, FL. February, 1984.

Training Course on Toxicology and Risk Assessment. Conducted for the Florida Department of Health and Rehabilitative Services. Orlando, Florida. October 3-5, 1984.

Monitoring and Management of Hazardous Waste at RCRA Subtitle D Facilities. Conducted for U.S. EPA Office of Research & Development. Tallahassee, Florida. October, 1984.

Symposium on EDB. Conducted for FL State University System. Tallahassee, FL. February, 1984.

Workshop on Monitoring Considerations in the Siting and Operation of Hazardous Waste Disposal Facilities in Temperate Zone Wet Environments. Conducted for the Office of Research and Development, U.S. Environmental Protection Agency. Tallahassee, Florida. October 4-5, 1983.

First National Conference on Waste Exchange and Resource Reuse. Tallahassee, Florida. March, 1983.

Training Course on Management of Hazardous Waste and Hazardous Materials. Conducted for the Florida Department of Environmental Regulation. Orlando, Florida. January 14-15, 1981.

Hazardous Materials Contingency Planning Workshop. Conducted for Region IV, Environmental Emergencies Branch, U.S. Environmental Protection Agency. Atlanta, Georgia. March 26, 1981.

Workshop on Hazardous Waste Management for Educational Institutions. Conducted for the Florida Department of Education. Orlando, Florida. February 3-4, 1981.

Hazardous Materials Incidents Workshop for Leon County, Florida. Conducted for the Leon County Commission. Tallahassee, Florida. July 14, 1980.

State Waste Exchange Workshop. For Florida Chamber of Commerce. Tallahassee, FL. December, 1980.

Hazardous Waste Management Seminar for Engineers. Conducted for the Florida Engineering Society. Tampa, FL. November, 1980.

ATTACHMENT C

**Testimony of Dr. Christopher M. Teaf
(May 2004 to May 2008)**

**Testimony of Dr. Christopher M. Teaf
(Period 5/04 through 5/08)**

CASE IDENTIFIER	JURISDICTION	STATE	TESTIMONY DATE	
			Deposition	Hearing/Trial
Broward County vs Southern Waste Systems and Sun Recycling	Broward County Administrative Hearing	FL	7/07	8/07
George vs Poseys on the River, Inc.	Second Judicial Circuit	FL	9/07	
Williams et al. vs City of Jacksonville et al.	Circuit Court for Duval County	FL	5/04; 6/04	
Morris et al. vs. McKibbin	Circuit Court for Hillsborough County	FL	4/05	
Wyatt vs. Chubb Group et al.	33rd Judicial District Court, Allen Parish	LA	3/08	3/08
Halley vs. Exxon et al.	Civil District Court, Jefferson Parish	LA		11/06
Peterson, et al., v. D.R. Horton, Inc.	Circuit Court for Montgomery County	MD	11/07	3/08
State of Oklahoma vs Tyson Foods et al.	U.S. District Court, Northern District	OK	1/08	2/08
Ponca Tribe et al. vs. Continental Carbon Company et al	U.S. District Court, Western District	OK	5/06	
Collins vs South Carolina Central RR et al.	Fourth Judicial Circuit, Williamsburg County	SC	10/05	
Watson vs Valley Proteins	U.S. District Court, Charleston Division	SC	9/04	

ATTACHMENT D

Tables & Figures

Table B1
Beneficial Use Designations for Select Waterbodies

Waterbody name	Waterbody ID Numbers	Fish & Wildlife Propagation beneficial use	Agriculture beneficial use	Recreation beneficial use	Aesthetics beneficial use	Limitations	Remarks
Upper Illinois River upstream of Baron Fork Confluence	121700030010, 121700030350, 121700030080	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	Scenic River
Tahlequah Creek (Town Branch)	121700030040	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	
Baron Fork from mouth upstream	121700050010	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	Scenic River
Tyner Creek	121700050090	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	
Peach eater Creek	121700050120	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	
Flint Creek	121700060010	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	Scenic River
Sager Creek	121700060080	Cool Water Aquatic Community	Yes	Primary Body Contact Recreation	Yes	Outstanding Resource Waters	

Table B2 ILLINOIS RIVER WATERSHED BACTERIAL SYNOPSIS							
	Exceedances of Geometric Mean Criterion (Recreational Season: 2005-2007)				Exceedances of Single Point Criterion (Recreational Season: 2005-2007)		
Illinois River Stream Reach	Parameter (cfu/100 mL)	GM	Dates	Number of Samples in Dataset (Data sources)	Parameter (cfu/100 mL)	Percent Exceedance (exceedance/total)	Magnitude of exceedances
OK121700060080- Sager Creek (4.2 miles)	Enterococci (33)	53	7/6/05-8/2/05	5 (CDM, OWRB)	E. coli (406)	4% (1/28)	One over 49x the std
	Enterococci (33)	81	8/16/05-9/13/05	7 (CDM, OWRB)	Enterococci (406)	11% (4/35)	Three over 3x the std; 1 over 60x the std
	Enterococci (33)	69	7/25/06-8/16/06	5 (CDM, OWRB)	Fecal coliform (400)	22% (8/36)	Three over 3x the std
OK121700060010- Flint Creek (7.8 miles)	Enterococci (33)	210 (6x)	6/29/06-7/25/06	8 (CDM, OWRB, USGS)	E. coli (235)	11.6% (11/95)	Three over 10x the std; one over 22x std
	Enterococci (33)	432 (13x)	8/1/06-8/23/06	6 (CDM, OWRB, USGS)	Enterococci (61)	65.5% (57/87)	Fourteen over 3x the std
	Enterococci (33)	158	5/2/06-5/31/06	5 (OCC)	Fecal coliform (400)	17% (7/42)	Five over 3x the std; one over 13x std
	Enterococci (33)	77	6/6/06-7/5/06	5 (OCC)			
	Enterococci (33)	164	7/11/06-8/7/06	5 (OCC)			
	Enterococci (33)	71	5/1/07-5/29/07	5 (OCC)			
	Enterococci (33)	202 (6x)	7/9/07-8/7/07	5 (OCC)			
	Enterococci (33)	190	8/20/07-9/18/07	5 (OCC)			
OK121700050010- Baron Fork (23.3 miles)	E. coli (126)	355	5/1/06-5/15/06	5 (CDM, USGS)	E. coli (235)	12% (6/51)	Two over 4x the std
	Enterococci (33)	981 (30x)	5/1/06-5/15/06	5 (CDM, USGS)	Enterococci (61)	30% (16/53)	Ten over 3x the std; seven over 6x the std
	Enterococci (33)	34	7/10/06-8/10/06	11 (CDM, OWRB, USGS)	Fecal coliform (400)	11% (6/53)	Two over 4x the std
	Enterococci (33)	36	8/16/06-8/29/06	5 (CDM, OWRB)			
	Enterococci (33)	70	9/18/06-9/27/06	5 (CDM, OWRB)			
	Fecal coliform (400)	439	5/1/06-5/15/06	5 (CDM, USGS)			
OK121700030010- Illinois River near Tahlequah (7.7 miles)	E. coli (126)	155	6/7/07-7/6/07	5 (USGS)	E. coli (235)	6% (3/47)	Two over 3x std
	Enterococci (33)	62	6/26/06-7/19/06	7 (CDM, OWRB)	Enterococci (61)	42% (21/50)	Eleven over 3x std; four over 6x std
	Enterococci (33)	49	8/1/06-8/29/06	9 (CDM, OWRB, USGS)	Fecal coliform (400)	10% (5/51)	Two over 4x std
	Enterococci (33)	33	8/29/06-9/27/06	5 (CDM, OWRB, USGS)			
	Enterococci (33)	180 (5x)	6/7/07-7/6/07	5 (USGS)			
OK121700030350- Illinois River near Watts (5.2 miles)	Enterococci (33)	62	8/1/06-8/24/06	5 (OWRB, USGS)	E. coli (235)	21% (6/28)	Four over 9x std
					Enterococci (61)	36% (12/33)	Six over 6x std; one at 125x std
					Fecal coliform (400)	19% (6/32)	Four over 11x std
OK121700030040- Tahlequah Creek (6.2 miles)		N/A			E. coli (406)	50% (2/4)	one over 29x std
					Enterococci (406)	100% (4/4)	three over 9x std; one over 29x std
					Fecal coliform (400)	100% (4/4)	four over 3x std; two over 30x std
OK121700030080- Illinois River: middle (31.5 miles)		N/A			E. coli (235)	15% (3/21)	two over 7x std; one over 17x std
					Enterococci (61)	36% (9/25)	six over 3x std; two over 65x std
					Fecal coliform (400)	15% (4/24)	three over 4x std; one over 19x the std
OK121700050090- Tyner Creek (15 miles)		N/A			E. coli (406)	22% (2/9)	two over 5x std
					Enterococci (406)	25% (3/12)	one over 2x std
					Fecal coliform (400)	45% (5/11)	three over 2x std; one over 11x std
OK121700050120- Peachwater Creek (10.3 miles)		N/A			E. coli (406)	17% (1/6)	
					Enterococci (406)	14% (1/7)	one over 3x std
					Fecal coliform (400)	29% (2/7)	one over 2x std; one over 4x std

CDM- Camp, Dresser & McKee; OWRB- Oklahoma Water Resources Board; USGS- United States Geological Survey; OCC-Oklahoma Conservation Commission

Table B3**NPDES Facilities in the IRW**

Facility name	Facility number	County	Receiving waters	SIC code	Permitted Flow (mgd)	Permitted Conc Average Limit (#/100mL)
OKLAHOMA						
Sequoyah Fuels Corporation	OK0000191	Sequoyah	Robert S. Kerr Reservoir (220200)	Industrial organic chemicals-2819	-	-
Souter Construction	OK0041106	Sequoyah	Illinois River	Crushed and broken limestone-1422	-	-
Stilwell Area Development Authority	OK0030341	Adair	Caney Creek (121700040010)	Sewerage system-4952	1.5	200
Tahlequah Public Works Authority	OK0026964	Cherokee	Tahlequah/IR (121700)	Sewerage system-4953	5.27	200
Westville Utility Authority	OK0028126	Adair	Shell/Baron Fork/TK (121700)	Sewerage system-4954	0.28	200

Source: EPA.GOV/ENVIRO/HTML/PCS/PCS_QUERY.HTML

Table B4

Estimated Fecal Coliform Sources in IRW (Oklahoma & Arkansas)

Estimated Total Fecal Coliform from All Sources by County (CFU/day)							
Counties in Watershed	Domestic Pets	Deer / Wildlife	Failing Septic Tanks	Point Sources	Livestock	Total	Percentage
Adair	1.30E+13	2.75E+12	2.15E+12	1.35E+10	2.10E+15	2.12E+15	18.1%
Benton	6.36E+12	1.24E+12	4.76E+13	1.70E+11	3.64E+15	3.70E+15	31.5%
Cherokee	1.98E+13	3.88E+12	2.80E+12	3.99E+10	5.18E+14	5.45E+14	4.6%
Delaware	3.48E+12	5.86E+11	7.05E+11	0.00E+00	4.21E+14	4.26E+14	3.6%
Sequoyah	3.89E+12	6.11E+11	5.05E+11	0.00E+00	1.10E+14	1.15E+14	1.0%
Washington	7.57E+12	2.26E+12	4.94E+13	6.15E+11	4.77E+15	4.83E+15	41.2%
Total	5.41E+13	1.13E+13	1.03E+14	8.38E+11	1.16E+16	1.17E+16	100.0%
Percentage	0.5%	0.1%	0.9%	0.01%	98.6%		100.0%

Estimated Total Fecal Coliform per Livestock Type by County (CFU/day)							
Counties in Watershed	Poultry (ALL)	Cattle & Calves	Horses & Ponies	Sheep & Lamb	Swine	Total	Percentage
Adair	4.15E+14	1.42E+15	6.97E+11	5.22E+12	2.55E+14	2.10E+15	18.1%
Benton	1.81E+15	1.27E+15	6.75E+11	3.95E+12	5.52E+14	3.64E+15	31.5%
Cherokee	4.68E+13	4.67E+14	3.88E+11	7.80E+11	2.55E+12	5.18E+14	4.5%
Delaware	1.12E+14	1.88E+14	1.11E+11	4.20E+11	1.20E+14	4.21E+14	3.6%
Sequoyah	5.43E+12	1.03E+14	1.26E+11	7.20E+10	8.77E+11	1.10E+14	0.9%
Washington	2.36E+15	1.68E+15	1.27E+12	5.02E+12	7.26E+14	4.77E+15	41.3%
Total	4.75E+15	5.13E+15	3.27E+12	1.55E+13	1.66E+15	1.16E+16	100.0%
Percentage	41.1%	44.4%	0.03%	0.1%	14.3%		100.0%

Table B5									
2002 (unless otherwise noted) USDA county agricultural census									
County	Cattle & Calves-all	Hogs & Pigs (Benton & Delaware 1997	Broilers	Turkeys (1997 for Adair & Delaware; 1978 for Sequoyah)	Horses & Ponies	Sheep & Lambs	Layers (1982 for Cherokee)	Pullets (1997 for Cherokee)	Poultry (broilers, layers
Numbers used	Sales only divided by 2	Inv + Sales divided by 2	Sales only divided by 5	Sales only divided by 2	Inv only	Sales only	Inv + Sales	Inv + Sales divided by 2.5	
Adair	17087	29536	2588549	154250	2077	544	949124	358736.4	3537673
Benton	27,086	113504	25,613,322	2073239	3570	731	2228954	901890	27842275.8
Cherokee	12591.5	663	718,801	262,327	2591	182	36575	70338	755376.2
Delaware	20125.5	123881	7430987	193301.5	2953	387	1411643	447482.8	8842630
Sequoyah	11726.5	959.5	205021	51250	3547	66	177181	110531.6	382202
Washington	26405.5	110140	21,978,106	1470501	4963	685	4865575	873153.6	26843681
Total	115022	378683.5	58534786	4204868	19701	2595	9669052	2762132.4	68203838

ANIMALS ON ANY GIVEN DAY

2002 (unless otherwise noted) USDA county agricultural census									
DENSITIES (animals/acre of pastureland in one year)									
County	Acres of pasture	Cattle & Calves-all	Hogs & Pigs	Poultry (broilers and layers	Horses & Ponies	Sheep & Lambs	Ducks and Geese	Pullets	Turkey
Adair	142120	0.1202	0.2078	24.8922	0.0146	0.0038		2.5242	1.0854
Benton	247726	0.1093	0.4582	112.3914	0.0144	0.0030		3.6407	8.3691
Cherokee	186520	0.0675	0.0036	4.0498	0.0139	0.0010		0.3771	1.4064
Delaware	219868	0.0915	0.5634	40.2179	0.0134	0.0018		2.0352	0.8792
Sequoyah	203569	0.0576	0.0047	1.8775	0.0174	0.0003		0.5430	0.2518
Washington	253091	0.1043	0.4352	106.0634	0.0196	0.0027		3.4500	5.8102
Adair and Cherokee average		0.0939	0.1057	14.4710	0.0143	0.0024		1.4506	1.2459

OBTAIN DENSITY BY DIVIDING ANIMALS/DAY BY PASTURE (ACRES)

2002 (unless otherwise noted) USDA county agricultural census									
Number of animals in one year									
	Acres of pasture	Cattle & Calves-all	Hogs & Pigs	Poultry (broilers and layers)	Horses & Ponies	Sheep & Lambs	Ducks and Geese	Pullets	Turkey
County in IRW									
Adair	113515	13648	23591	2825633	1659	435		286532	123204
Benton	111484	12189	51080	12529845	1607	329		405877	933019
Cherokee	66465	4487	236	269173	923	65		25064	93478
Delaware	19710	1804	11105	792695	265	35		40114	17328
Sequoyah	17232	993	81	32353	300	6		9356	4338
Washington	154380	16107	67183	16374061	3027	418		532605	896974
8-digit basin									
121700-03 (Adair & Cherokee)	53225	4996	5625	770219	759	128		77210	66312
121700-04 (Adair & Cherokee)	28587	2683	3021	413682	407	69		41470	35616
121700-05 (Adair & Cherokee)	74200	6965	7842	1073748	1058	178		107638	92445
121700-06 (Delaware)	15001	1373	8452	603309	201	26		30531	13188

DENSITY x PASTURE = IRW FIGURES

2002 (unless otherwise noted) USDA county agricultural census															
FC production / day	Cattle & Calves-all 1.04E+11 org / day		Hogs & Pigs 1.08E+10 org / day		Broilers and layers 1.36E+8 org / day		Horses & Ponies 4.20E+8 org / day		Sheep & Lambs 1.20E+10 org / day		Ducks and Geese 1.20E+10 org / day	Pullets 6.8E+7 org / day	Turkey 9.3E+7 org / day		
	Number of animals	Total FC (organisms / day)	Number of animals	Total FC (organisms / day)	Number of (organisms / day)	Total FC (organisms / day)	Number of (organisms / day)	Total FC (organisms / day)	Number of (organisms / day)	Total FC (organisms / day)	Number of (organisms / day)	Total FC (organisms / day)	Number of (organisms / day)		
County in IRW															
Adair	13648	1.42E+15	23591	2.55E+14	2825633	3.84E+14	1659	6.97E+11	435	5.22E+12		286532	1.9484E+13	123204	1.1125E+13
Benton	12189	1.27E+15	51080	5.52E+14	12529845	1.70E+15	1607	6.75E+11	329	3.95E+12		405877	2.76E+13	933019	8.4252E+13
Cherokee	4487	4.67E+14	236	2.55E+12	269173	3.66E+13	923	3.88E+11	65	7.80E+11		25064	1.7044E+12	93478	8.4411E+12
Delaware	1804	1.88E+14	11105	1.20E+14	792695	1.08E+14	265	1.11E+11	35	4.20E+11		40114	2.7278E+12	17328	1.5647E+12
Sequoyah	993	1.03E+14	81	8.77E+11	32353	4.40E+12	300	1.26E+11	6	7.20E+10		9356	6.3621E+11	4338	3.9172E+11
Washington	16107	1.68E+15	67183	7.26E+14	16374061	2.23E+15	3027	1.27E+12	418	5.02E+12		532605	3.6217E+13	896974	8.0997E+13
Total		5.12E+15		1.66E+15		4.46E+15		3.27E+12		1.55E+13			8.84E+13		1.87E+14
8-digit basin															
121700-03 (Adair & Cherokee)	4996	5.20E+14	5625	6.08E+13	770219	1.05E+14	759	3.19E+11	128	1.54E+12		77210	5.2503E+12	66312	5.988E+12
121700-04 (Adair & Cherokee)	2683	2.79E+14	3021	3.26E+13	413682	5.63E+13	407	1.71E+11	69	8.28E+11		41470	2.82E+12	35616	3.2161E+12
121700-05 (Adair & Cherokee)	6965	7.24E+14	7842	8.47E+13	1073748	1.46E+14	1058	4.44E+11	178	2.14E+12		107638	7.3194E+12	92445	8.3478E+12
121700-06 (Delaware)	1373	1.43E+14	8452	9.13E+13	603309	8.21E+13	201	8.44E+10	26	3.12E+11		30531	2.0761E+12	13188	1.1909E+12

Total FC/day 2002 USDA Census Data		
	Total IRW	
Cattle & Calves	5.12E+15	
Hogs & Pigs	1.66E+15	
Broilers & Layers	4.46E+15	
Horses & Ponies	3.27E+12	
Sheep & Lambs	1.55E+13	
Ducks & Geese		Have not been able to determine number of animals in counties
Pullets	8.84E+13	
Turkeys	1.87E+14	
Broilers, layers, pullets, turkeys	4.74E+15	Pullets estimated to release half as much FC as grown chickens
Cattle & Calves	5.12E+15	
All other livestock	6.79E+15	

SOURCES: EPA.GOV/MRLC/NLCD-2001.HTML, CANADIAN RIVER TMDL, 2002 USDA CENSUS

Note: FC Counts / day were obtained from the 1999 study by the American Society of Agricultural Engineers (ASAE)

Table B6
EDGE-OF-FIELD WATER SAMPLES EQUIVALENT TO RAW SEWAGE SPILLS

StationID	Collection Date	Bacteria	Bacterial Growth		
EOF01	5/23/05	E. coli		110,000	MPN* / 100ml
EOF01	5/23/05	Enterococcus Group	≥	1,600,000	MPN* / 100ml
EOF01	5/23/05	Fecal Coliform		110,000	MPN* / 100ml
EOF01	5/23/05	Total Coliform	≥	1,600,000	MPN* / 100ml
EOF-EOF1	6/17/06	Total Coliform		760,000	MPN* / 100ml
EOF02	5/23/05	Enterococcus Group		500,000	MPN* / 100ml
EOF02	5/23/05	Fecal Coliform		1,600,000	MPN* / 100ml
EOF02	5/23/05	Total Coliform	≥	1,600,000	MPN* / 100ml
EOF15	6/2/05	Total Coliform		160,000	MPN* / 100ml
EOF16	6/5/05	Total Coliform		170,000	MPN* / 100ml
EOF17	6/5/05	Total Coliform		220,000	MPN* / 100ml
EOF07	6/5/05	Total Coliform		110,000	MPN* / 100ml
EOF20	6/5/05	E. coli		300,000	MPN* / 100ml
EOF20	6/5/05	Enterococcus Group		300,000	MPN* / 100ml
EOF20	6/5/05	Fecal Coliform		300,000	MPN* / 100ml
EOF20	6/5/05	Total Coliform	≥	1,600,000	MPN* / 100ml
EOF18	6/5/05	Total Coliform		130,000	MPN* / 100ml
EOF-SPREAD023	4/25/06	E. coli		400,000	MPN* / 100ml
EOF-SPREAD023	4/25/06	Enterococcus Group		180,000	MPN* / 100ml
EOF-SPREAD023	4/25/06	Fecal Coliform		400,000	MPN* / 100ml
EOF-SPREAD023	4/25/06	Total Coliform		760,000	MPN* / 100ml
EOF-SPREAD029	4/25/06	Total Coliform		140,000	MPN* / 100ml
EOF-SPREAD007	4/25/06	Total Coliform		110,000	MPN* / 100ml
EOF-SPREAD060	4/29/06	Total Coliform		180,000	MPN* / 100ml
SPREAD053E	4/29/06	Enterococcus Group		270,000	MPN* / 100ml
SPREAD017A	5/1/06	E. coli		760,000	MPN* / 100ml
SPREAD017A	5/1/06	Enterococcus Group		1,200,000	MPN* / 100ml
SPREAD017A	5/1/06	Fecal Coliform		760,000	MPN* / 100ml
SPREAD017A	5/1/06	Total Coliform		1,200,000	MPN* / 100ml
SPREAD053G	5/4/06	Enterococcus Group		180,000	MPN* / 100ml
SPREAD053G	5/4/06	Total Coliform		540,000	MPN* / 100ml
EOF-SPREAD064	5/4/06	E. coli		540,000	MPN* / 100ml
EOF-SPREAD064	5/4/06	Enterococcus Group		1,200,000	MPN* / 100ml
EOF-SPREAD064	5/4/06	Fecal Coliform		540,000	MPN* / 100ml
EOF-SPREAD064	5/4/06	Total Coliform		760,000	MPN* / 100ml
EOF-SPREAD048	5/9/06	E. coli		140,000	MPN* / 100ml
EOF-SPREAD048	5/9/06	Enterococcus Group		180,000	MPN* / 100ml
EOF-SPREAD048	5/9/06	Fecal Coliform		140,000	MPN* / 100ml
EOF-SPREAD048	5/9/06	Total Coliform		400,000	MPN* / 100ml
EOF-SPREAD010	5/9/06	E. coli		180,000	MPN* / 100ml
EOF-SPREAD010	5/9/06	Enterococcus Group		540,000	MPN* / 100ml
EOF-SPREAD010	5/9/06	Fecal Coliform		180,000	MPN* / 100ml
EOF-SPREAD010	5/9/06	Total Coliform	>	1,200,000	MPN* / 100ml
EOF-SPREAD071	5/10/06	Enterococcus Group		140,000	MPN* / 100ml
SPREAD073B	6/18/06	E. coli		1,200,000	MPN* / 100ml
SPREAD073B	6/18/06	Enterococcus Group	>	1,200,000	MPN* / 100ml
SPREAD073B	6/18/06	Fecal Coliform		1,200,000	MPN* / 100ml
SPREAD073B	6/18/06	Total Coliform	>	1,200,000	MPN* / 100ml
EOF-SPREAD023	6/18/06	Fecal Coliform		180,000	MPN* / 100ml
EOF-SPREAD023	6/18/06	Total Coliform		760,000	MPN* / 100ml

TABLE T1
Disinfection Byproducts in IRW Water Supplies
(# of exceedances/total # of samples)

Public Water System using IRW waters as a source	Chloroform		Dibromochloromethane	Bromodichloromethane	TTHM	HAA5
	MCLG (70 µg/L)	Risk-based ^a (0.17 µg/L)	Risk-based ^b (0.9 µg/L)	Risk-based ^b (1 µg/L)	MCL (80 µg/L)	MCL (60 µg/L)
Adair Co. RWD#5	0/18	12/18	8/16	11/16	0/16	0/11
Burnt Cabin RWD	4/16	16/16	3/11	8/11	3/12	2/9
Cherokee Co. RWD #2	9/51	50/51	39/42	42/42	9/42	2/18
Cherokee Co. RWD #13	5/34	34/34	10/24	23/24	4/26	1/15
East Central OK Water Authority	2/44	43/44	21/30	30/30	4/30	5/16
Fin & Feather Resort	0/18	17/18	7/8	8/8	0/8	0/5
Flintridge POA	0/22	15/22	16/17	17/17	0/17	0/10
Gore PWA	1/18	17/18	4/15	15/15	1/18	7/15
LRed (Chicken Creek)	1/23	21/23	0/14	6/14	1/14	0/10
LRed (Lakewood)	1/16	11/16	1/9	5/9	1/9	1/8
LRed (Wildcat)	1/19	18/19	4/10	6/10	0/11	1/6
LRed (Woodhaven)	4/24	24/24	5/15	15/15	3/16	3/10
Pettit Mt. Water	1/20	19/20	5/7	7/7	1/7	0/2
Sequoyah Co. RWD #5	2/28	28/28	21/22	22/22	2/22	4/17
Sequoyah County WA	10/72	71/72	25/60	60/60	8/64	0/36
Tahlequah PWA	4/345	297/345	62/133	124/133	10/148	6/107
Tenkiller Aqua Park	2/25	23/25	12/13	13/13	4/13	0/4
Tenkiller Utility Co.	4/23	22/23	15/19	18/19	5/19	3/14
Total	51/816	738/816	258/465	430/465	56/492	35/313
% Exceedances	6.3%	90.4%	55.5%	92.5%	11.4%	11.2%

^a U.S. EPA Region 6 health-based screening level

^b Risk-based values listed in Stage 2 DBP Rule

MCL and MCLG as listed in Stage 2 DBP Rule

Source: Oklahoma Department of Environmental Quality

TTHM-Total Trihalomethanes

HAA5- Haloacetic acids

IRW- Illinois River Watershed

TABLE T2
Disinfection Byproducts in Selected IRW Water Supplies
(# of exceedances/total # of samples)

LOCATION	Chloroform (µg/L)		Bromodichloromethane (µg/L)		Dibromo- chloromethane (µg/L)	Dichloroacetic acid (µg/L)	Trichloroacetic acid (µg/L)	TTHM (µg/L)	HAA5 (µg/L)
	70 MCLG	0.17 Screening level	0.18 Screening level	1 Risk based	0.9 Risk based	0.7 Risk based	20 MCLG	80 MCL	60 MCL
Cherokee RWD #2									
Treatment Plant	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Jimmy Houston	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
John Bates	1/3	3/3	3/3	3/3	3/3	3/3	0/3	2/3	0/3
Keys Fire	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Maurice	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Gore PWA									
ACOE Office	0/3	3/3	3/3	3/3	3/3	3/3	1/3	0/3	0/3
City Complex	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Gore HS	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Marval Trout Resort	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Treatment Plant	0/3	3/3	3/3	3/3	3/3	3/3	0/3	0/3	0/3
Tahlequah									
Dano's	0/3	3/3	3/3	3/3	3/3	3/3	1/3	0/3	0/3
EZ-Mart	0/3	3/3	3/3	3/3	3/3	3/3	1/3	1/3	1/3
Treatment Plant	0/3	3/3	3/3	3/3	3/3	3/3	1/3	0/3	1/3
Wellington Store	0/3	3/3	3/3	3/3	3/3	3/3	1/3	0/3	0/3
WPC Plant	1/3	3/3	3/3	3/3	3/3	3/3	1/3	3/3	1/3
Ratio of Exceedances	2/45	45/45	45/45	45/45	45/45	45/45	6/45	6/45	3/45
Percent Exceedances	4.4 %	100 %	100 %	100 %	100 %	100 %	13.3 %	13.3 %	6.7 %

Source: CDM, 2007

TABLE T3
Trihalomethane Forming Potential (THMFP)
 May 2005-August 2007

Average THMFP reported as CHCl_3 ($\mu\text{g/L}$)
 (ranges)

Tahlequah	Cherokee Co. RWD #2	Cherokee Co. RWD #13	Gore PWA	Adair Co. RWD #5
84.5* (32.4-184.4)	165.1* (20.7-405.4)	149.5* (18.3-228.3)	135.1* (14-221.8)	59.3 (21.8-136.1)

* Exceeds EPA drinking water standard of 80 $\mu\text{g/L}$ for Total Trihalomethanes

Table C1. Lake Tenkiller Chlorophyll a (µg/L)				
Date	Station			
	LK-01	LK-02	LK-03	LK-04
5/18/05	No data	11.2	28.6	No data
6/2/05	15.5	12.2	11.4	No data
6/15/05	9.4	14.5	26.8	No data
6/29/05	4.1	5.8	14	No data
7/12/05	8.8	10.2	15.8	No data
7/26/05	4.2	12	19.1	36.1
8/9/05	9.5	13.8	9.7	14.9
8/23/05	11.4	11.8	12.5	62.8
9/7/05	10.2	10.6	9.9	17
9/20/05	7.6	9.4	8.3	133.3
10/4/05	4.3	7.3	12.8	42.6
11/16/05	6.4	5.3	9.5	0.6
3/28/06	1.4	1.7	45.8	0.6
4/6/06	3.4	0.7	16.7	1.6
4/18/06	0.6	1.1	11.5	9.5
5/2/06	1.1	2	5.3	1.2
5/16/06	2.1	12.8	20	9.8
6/1/06	5	17	19.7	20.2
6/14/06	4.1	9.8	12.5	26.2
6/27/06	4.6	9.3	15.7	39.7
7/13/06	4.7	6.1	5.8	24.4
6/12/07	14.6	11.7	24.4	26.1
7/17/07	6.2	6.8	16.3	22.9
8/29/07	7.8	5.4	10.2	24.6
Average	6.4	8.7	15.9	27.1
Source: CDM, 2008				

TABLE C2 Exceedances of WHO Guidelines on Lake Tenkiller Reservoir*		
WHO Guidance	Cells/mL	Lake Tenkiller
Low probability of adverse health effects	20,000	58% ^a
Moderate probability of adverse health effects	100,000	24% ^b
High probability of adverse health effects	Visible scums	One sample greater than 1,000,000 cells/mL

*OWRB 2004, 2005; CDM 2006, 2007

^aPercentage of samples greater than 20,000 cells/mL

^bPercentage of samples greater than 20,000 that are also greater than 100,000 cells/mL

TABLE C3
Army Corps of Engineers Data

Location	Total Cyanobacteria expressed as Biovolume ($\mu\text{m}^3/\text{mL}$)										
	4/26/01	5/24/01	6/21/01	7/26/01	8/16/01	9/20/01	10/18/01	11/15/01	1/28/02	3/14/02	4/11/02
Dam	253,688	41,245	8,714,700	10,227,215	23,046,105	6,311,574	535,773	367,880	30,381	27,140	No data
Buckhorn	No data	309,162	14,386,980	13,450,149	50,203,158	10,874,913	1,695,290	62,500	97,009	345,334	106,312
Carlile	255,818	1,725,624	13,307,413	41,444,830	70,874,631	4,902,506	2,767,424	798,375	130,724	119,989	21,163
Cherokee	4,840	551,682	16,830,542	53,168,181	15,354,976	2,726,043	2,865,366	571,542	12,559,751	69,779	38,763
Location	5/9/02	6/12/02	7/10/02	8/8/02	9/13/02	10/17/02	11/14/02	1/28/03	3/13/03	4/17/03	5/6/03
Dam	267,769	11,595,678	35,319,545	60,231,258	18,827,278	6,009,327	129,864	595,551	838,341	4,692	30,931
Buckhorn	33,889	10,811,846	40,637,254	61,208,057	41,480,143	7,003,648	1,424,175	450,939	1,095,037	35,632	72,850
Carlile	498,294	13,663,420	47,116,087	54,752,131	41,137,780	1,995,079	681,046	303,610	158,971	78,450	629,569
Cherokee	1,098,711	12,664,686	No data	14,011,676	35,633,410	9,139,743	50,819	37,813	35,437	1,171,449	4,487,341
Location	6/1/03	7/8/03	8/4/03	9/8/03	10/9/03	11/13/03	5/12/04	6/16/04	7/14/04	8/12/04	9/20/04
Dam	216,294,706	146,539,063	556,117,872	296,091,837	167,799,435	95,396,609	377,696	6,925,795,238	2,493,446,100	7,530,096,500	1,883,080,000
Buckhorn	197,986	1,381,561,856	421,182,708	2,164,205,239	572,656,250	906,832	319,520,000	6,143,000,000	8,455,895,238	2,156,877,900	5,236,800,000
Carlile	78,045,806	2,712,078,947	6,569,702,381	1,487,762,500	756,706,206	1,239,297,354	No data	3,891,750,000	9,695,095,238	3,304,300,000	8,454,024,900
Cherokee	351,122	3,886,962,206	3,937,724,311	60,802,075,714	4,009,435,897	896,374	425,973,600	7,944,580,800	10,335,098,300	5,710,053,600	10,894,444,500

Source: ACoE





Figure B2

EPA's Logarithmic Exposure – Response Model



Adapted from USEPA Implementation Guidance for Ambient Water Quality Criteria for Bacteria (2003)

Figure B3. Land Application of Poultry Waste Vs. Illinois River Recreation Use




Figure B4. Rainfall vs. River Floaters vs. Monthly % Land Application





Figure B5







Figure B7



Well Sampling Locations
In the Illinois River Watershed


Sampling Sites

- ▲ Wells

▲ Wells

Geoprobe Sampling Locations
In the Illinois River Watershed

Figure B8



Sampling Sites

• Geoprobe






Figure B10**REPORTED CAMPYLOBACTER INFECTION RATE**

Figure B11



REPORTED SALMONELLA INFECTION RATE

Figure T1

THMs/HAA5s Related To Eutrophication

Poultry Waste Disposal
High Nutrient Levels




Runoff

Source Water Eutrophication
Algal Growth



Withdrawal



Public
Water
Supply

Chlorine
Disinfection

THMs
(potential carcinogens)



Eighteen Currently Active Public Water Supply
Systems Whose Source is Within the IRW






Figure T2





Army Corps of Engineers □
Cyanobacteria Sampling Locations




Figure C2

■ Corps of Engineers Stations